



A Review on the Potential of Ginger Essential Oil-based Nanotechnology for Controlling Tropical Plant Diseases

**Taheera Ansari ^{a++*}, Sudhakar S ^{b#},
Narsing Laxmi Prasanna ^{c++}, Rashmi Nigam ^{d†},
Uma Shankar ^{e‡}, Deepali Mohapatra ^{f^},
Lipikant Sahoo ^{f^} and S. Syam Prasad ^{g##}**

^a Department of Zoology, Deen Dayal Upadhyaya Gorakhpur University, Gorakhpur, Uttar Pradesh- 273009, India.

^b Plant Protection, KVK, IDUKKI, India.

^c Department of Plant Pathology, Navsari Agricultural University, Navsari- 396 450, India.

^d Department of Plant Pathology, J.V. College Baraut (Baghpat), C.C.S. University Meerut, Uttar Pradesh-250611, India.

^e Department of Plant Pathology, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya, India.

^f Department of Plant Pathology, Odisha University of Agriculture and Technology (OUAT), Bhubaneswar, Odisha, India.

^g Department of Botany, Government Degree College, Nellakondapally, Telangana, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2025/v28i11876>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129015>

⁺⁺ Ph.D. Scholar;

[#] Scientist;

[†] Associate Professor;

[‡] Ph.D;

[^] Ph.D. Research Scholar;

^{##} Assistant Professor;

*Corresponding author: E-mail: ansaritaheera@gmail.com;

Cite as: Ansari, Taheera, Sudhakar S, Narsing Laxmi Prasanna, Rashmi Nigam, Uma Shankar, Deepali Mohapatra, Lipikant Sahoo, and S. Syam Prasad. 2025. "A Review on the Potential of Ginger Essential Oil-Based Nanotechnology for Controlling Tropical Plant Diseases". *Journal of Advances in Biology & Biotechnology* 28 (1):230-48. <https://doi.org/10.9734/jabb/2025/v28i11876>.

ABSTRACT

The increasing prevalence of plant diseases in tropical agriculture, driven by climate change, pathogen resistance, and unsustainable pesticide use, necessitates the development of eco-friendly and effective disease management strategies. Ginger essential oil (GEO), derived from *Zingiber officinale*, is a natural antimicrobial agent with proven efficacy against a wide range of plant pathogens, including bacteria, fungi, and viruses. Its bioactive compounds, such as gingerols, shogaols, and zingiberene, exhibit multiple modes of action, making GEO a promising alternative to synthetic pesticides. Its volatility, short shelf life, and limited solubility under field conditions restrict its direct application in agriculture. Nanotechnology offers a transformative solution to these limitations by encapsulating GEO in advanced nanocarriers, such as liposomes, nanoemulsions, and polymeric nanoparticles, which improve its stability, bioavailability, and controlled release. Nanoformulated GEO has demonstrated enhanced antimicrobial efficacy in both laboratory and field studies, outperforming pure GEO and synthetic agrochemicals in controlling plant pathogens like *Fusarium oxysporum*, *Alternaria alternata*, and *Xanthomonas campestris*. GEO nanoformulations are biodegradable and pose minimal risks to non-target organisms and the environment, aligning with the principles of sustainable agriculture. Despite these promising outcomes, challenges such as high production costs, regulatory uncertainties, and limited farmer awareness hinder the large-scale implementation of GEO nanotechnology. It should focus on optimizing nanoformulation techniques, exploring synergistic combinations with other natural antimicrobials, and integrating GEO nanoformulations into integrated pest management (IPM) strategies. Collaboration between researchers, policymakers, and farmers is essential to overcome adoption barriers and ensure effective deployment. By addressing these challenges, GEO-based nanotechnology has the potential to revolutionize tropical agriculture, providing a sustainable, safe, and innovative solution for managing plant diseases while reducing dependency on harmful chemical pesticides.

Keywords: Nanotechnology; plant disease management; nanoformulations.

1. INTRODUCTION

A. Plant Disease Management

Tropical plant diseases are a significant threat to global agriculture, with devastating effects on crop production, food security, and farmer livelihoods, particularly in developing countries. These diseases, caused by fungi, bacteria, viruses, nematodes, and other pathogens, thrive in warm and humid tropical environments, leading to severe epidemics in key crops such as rice, banana, cassava, cocoa, and coffee (Jones *et.al.*, 2012). For example, diseases like Fusarium wilt in bananas and Black Sigatoka reduce banana yields by 40-50%, threatening a crop that is a staple food for millions in tropical regions. Similarly, rice blast disease, caused by *Magnaporthe oryzae*, destroys up to 30% of global rice production annually, directly impacting global food supplies. The expansion of global trade, monoculture farming practices, and

climate change exacerbate the spread and severity of these diseases. The introduction of invasive pathogens into non-native regions has created emerging disease hotspots, increasing the challenge of containment and control (Carnegie *et.al.*, 2019). This underscores the critical need for effective disease management strategies to mitigate the economic and ecological damage caused by tropical plant pathogens. The economic burden of tropical plant diseases is immense. Annually, crop losses due to pathogens exceed \$220 billion globally, with tropical regions bearing the highest share of these losses. For smallholder farmers, who constitute the majority of agricultural producers in tropical areas, these losses can have catastrophic consequences, perpetuating cycles of poverty and food insecurity. The downstream economic impacts, including increased food prices and reduced export revenues, strain national economies and disrupt international trade. From an ecological perspective, plant

pathogens significantly affect biodiversity and ecosystem services. Diseases like coffee leaf rust (*Hemileia vastatrix*) not only reduce coffee yields but also threaten shade-grown coffee plantations that support diverse ecological systems (Avelino *et al.*, 2023). The indiscriminate use of chemical pesticides to combat plant diseases contributes to soil degradation, water contamination, and harm to non-target organisms, disrupting delicate ecological balances. Thus, addressing tropical plant diseases is crucial not only for agricultural sustainability but also for preserving environmental health.

B. Conventional Disease Management Strategies

Chemical pesticides have long been the cornerstone of plant disease management due to their rapid and broad-spectrum action. Their widespread and often indiscriminate use has led to numerous limitations, including the development of pathogen resistance. Fungicides like azoles and carbamates, widely used to control fungal diseases, are increasingly ineffective due to the evolution of resistant strains of pathogens such as *Botrytis cinerea* and *Fusarium oxysporum* (Oliver *et.al.*, 2014). The overuse of chemical pesticides has severe environmental consequences, including the contamination of soil and water resources, bioaccumulation in food chains, and harm to beneficial organisms like pollinators and soil microbes. Human health concerns associated with pesticide exposure further highlight the limitations of chemical approaches. Chronic exposure to pesticides has been linked to respiratory illnesses, cancers, and neurological disorders in farming communities. Consequently, regulatory restrictions on pesticide use and growing consumer demand for pesticide-free produce have prompted the search for alternative, sustainable disease management strategies. Biological control methods, which leverage natural antagonists such as beneficial microbes, predatory insects, and natural products, have emerged as eco-friendly alternatives to chemical pesticides. For example, *Trichoderma spp.* are widely used as biocontrol agents against soilborne pathogens due to their ability to outcompete pathogens and induce plant resistance. Similarly, bacterial species like *Bacillus subtilis* and *Pseudomonas fluorescens* have shown promise in suppressing fungal and bacterial diseases in crops (Santoyo *et.al.*, 2012). Despite their potential, biological control

methods face several challenges. Their efficacy can be inconsistent due to environmental variables such as temperature, humidity, and soil pH, which affect the survival and activity of biocontrol agents. Scaling up the production and commercialization of biocontrol products remains a significant hurdle, limiting their widespread adoption in tropical agriculture. Therefore, while biological control methods are valuable components of integrated disease management strategies, they require further refinement and supplementation with other innovative approaches.

C. Need for Novel Approaches in Plant Disease Control

Given the limitations of conventional strategies, there is an urgent need for sustainable and eco-friendly approaches to plant disease management. Such approaches prioritize minimizing environmental harm, reducing pesticide residues in food, and promoting agricultural sustainability. Plant-derived natural products, particularly essential oils, have garnered attention as promising alternatives due to their antimicrobial, antifungal, and antiviral properties (Wani *et.al.*, 2021). Unlike synthetic pesticides, essential oils are biodegradable, pose minimal risks to human and animal health, and are less likely to induce resistance in pathogens. Sustainable strategies like integrated pest management (IPM), which combine biological, cultural, and chemical controls, are increasingly recognized as essential for addressing the multifaceted challenges of tropical plant disease management. These approaches must be complemented by innovative technologies to enhance their efficacy and scalability. Nanotechnology, which involves manipulating materials at the nanometer scale, offers transformative potential for agriculture, including plant disease management. By engineering nanoparticles and nanoformulations, researchers can enhance the delivery, stability, and bioavailability of active compounds such as essential oils. Nanoemulsions and nanocapsules can protect volatile compounds in essential oils from degradation, ensuring prolonged efficacy against plant pathogens. Nanotechnology enables targeted delivery of antimicrobial agents, reducing off-target effects and environmental contamination. The application of nanotechnology in agriculture extends beyond pest control, encompassing areas such as precision farming, nutrient delivery, and pathogen detection (Yadav *et al.*, 2023). By integrating nanotechnology with

natural products like ginger essential oil (GEO), it is possible to develop innovative, sustainable solutions for combating tropical plant diseases while addressing the limitations of existing approaches.

D. Objectives and Scope of the Review

The primary objective of this review is to explore the potential of ginger essential oil (GEO), derived from *Zingiber officinale*, as a natural antimicrobial agent for managing tropical plant diseases. GEO is rich in bioactive compounds, including gingerols, shogaols, and zingiberene, which exhibit potent antimicrobial and antifungal properties. Despite its promising bioactivity, the volatility and instability of GEO in field applications necessitate innovative delivery systems to enhance its efficacy and durability. This review also examines the integration of nanotechnology with GEO to develop advanced formulations capable of overcoming its limitations. By encapsulating GEO in nanocarriers such as liposomes, nanoparticles, and nanoemulsions, it is possible to improve its stability, control its release, and enhance its pathogen-targeting capabilities (Rashwan *et.al.*, 2023). The scope of this review includes an evaluation of existing studies on GEO-based nanotechnology, its potential for application in tropical agriculture, and the challenges and opportunities for scaling up these technologies.

2. TROPICAL PLANT DISEASES: A GROWING CONCERN

A. Major Tropical Plant Diseases

Fungal diseases represent some of the most destructive threats to tropical crops, with *Fusarium* species and *Mycosphaerellafijiensis* among the leading pathogens. Fusarium wilt, caused by *Fusarium oxysporum* species complex, is a globally significant fungal disease that targets crops such as bananas, tomatoes, and legumes. Tropical agriculture is particularly vulnerable to the *Fusarium* Tropical Race 4 (TR4) strain, which affects Cavendish bananas and has spread widely across Southeast Asia, Africa, and Latin America. This disease invades the vascular system of plants, causing wilting, stunted growth, and eventual death, leading to severe yield losses of up to 100% in infected plantations (Elangovan *et al.*, 2024). Its resilience to chemical control and capacity to persist in soil for decades exacerbate its threat. Another major

fungal disease in tropical agriculture is Black Sigatoka, caused by *Mycosphaerellafijiensis*. This foliar disease severely impacts banana and plantain production by reducing photosynthetic leaf area, leading to premature fruit ripening and yield losses of 35-50% if left untreated. Black Sigatoka thrives in the humid climates of tropical regions, and its management typically requires intensive fungicide use. The rapid emergence of fungicide-resistant strains of *M. fijiensis* complicates control efforts and underscores the need for innovative management strategies. Bacterial diseases also pose significant challenges to tropical agriculture, with bacterial blight of rice being one of the most devastating examples. This disease is caused by *Xanthomonas oryzae* *pv.* *oryzae* (Xoo) and affects rice, a staple crop for over half of the world's population. Bacterial blight thrives in warm, humid conditions typical of tropical regions, making rice fields in Southeast Asia, Africa, and Latin America highly vulnerable (Khalaf *et.al.*, 2024). The pathogen invades rice plants through natural openings or wounds, causing leaf yellowing, wilting, and death. Infected fields can suffer yield losses ranging from 20% to 80%, depending on disease severity. Efforts to manage bacterial blight often rely on resistant rice varieties; the emergence of virulent pathogen strains capable of overcoming host resistance genes has rendered many traditional approaches ineffective. Chemical controls for bacterial diseases remain limited, leaving farmers with few effective tools for managing this pathogen. Viral diseases are a growing concern in tropical agriculture due to their ability to rapidly spread through insect vectors and infected plant material. Banana bunchy top virus (BBTV) is one of the most economically damaging viral diseases affecting banana crops in tropical and subtropical regions. Transmitted by the banana aphid (*Pentalonianigranervosa*), BBTV causes severe stunting, leaf distortion, and fruit yield losses of up to 100% in infected plants. BBTV has become increasingly difficult to manage due to the widespread presence of its aphid vector and the limited availability of virus-free planting materials. Cassava mosaic disease (CMD), caused by cassava mosaic geminiviruses (CMGs), is a major constraint on cassava production in tropical Africa and Asia. CMD is transmitted by whiteflies (*Bemisiatabaci*) and through infected stem cuttings, leading to leaf curling, chlorosis, and yield losses of up to 90% in severe cases (Bhaargavi *et.al.*, 2024). The disease not only reduces food security in regions where cassava

is a staple crop but also limits the crop's potential as an industrial bioresource.

B. Current Challenges in Disease Management

Climate change is a major driver of the increased prevalence and severity of plant diseases in tropical regions. Rising temperatures, changing precipitation patterns, and increased frequency of extreme weather events create conditions that favor pathogen proliferation and expand the geographic range of diseases. For example, warmer temperatures and high humidity enhance the growth and spread of fungal pathogens such as *Mycosphaerellafijiensis* (Black Sigatoka) and *Colletotrichum gloeosporioides* (anthracnose), leading to higher disease incidences in tropical crops. Climate change facilitates the expansion of pest and vector populations, such as whiteflies and aphids, which transmit viral diseases like CMD and BBTv, respectively (Mangang *et.al.*, 2024). The interaction between climate change and plant diseases is further compounded by the increased susceptibility of stressed plants to infections. Prolonged droughts, flooding, and soil salinity weaken plant immune responses, making crops more vulnerable to pathogens. These climate-induced challenges highlight the need for adaptive and resilient disease management strategies tailored to changing environmental conditions. Smallholder farmers dominate tropical agriculture, accounting for up to 80% of food production in regions like sub-Saharan Africa and Southeast Asia. These farmers often lack access to effective plant disease management solutions, leaving their crops highly vulnerable to pathogens. High costs of chemical pesticides and fungicides, coupled with inadequate availability of biocontrol agents and resistant crop varieties, limit the ability of smallholder farmers to protect their crops (Bardin *et.al.*, 2015). Limited infrastructure for pathogen diagnosis and early warning systems makes it difficult for farmers to detect and respond to disease outbreaks in a timely manner. Compounding these issues is the lack of farmer education and training in sustainable agricultural practices. Many smallholders continue to rely on traditional farming methods, which may exacerbate disease pressure through practices such as monocropping and improper pesticide application. Bridging these gaps in resources and knowledge is critical for empowering smallholder farmers to effectively manage tropical plant diseases. The overreliance on conventional pesticides has led to the

rapid emergence of pesticide-resistant pathogens and pests in tropical agriculture. For example, fungicide-resistant strains of *Mycosphaerellafijiensis* (Black Sigatoka) have emerged in banana-growing regions, rendering many fungicides ineffective and necessitating higher application rates, which increase costs and environmental risks. Similarly, bacterial pathogens like *Xanthomonas spp.* have developed resistance to commonly used antibiotics, further complicating the management of diseases such as bacterial blight of rice (Sharma *et.al.*, 2017). Pesticide resistance not only reduces the efficacy of disease control measures but also threatens the sustainability of tropical agriculture by promoting the overuse of chemicals. This results in environmental contamination, harm to non-target organisms, and disruption of ecological balances. Developing integrated pest and disease management strategies that reduce dependency on chemical pesticides is essential to overcoming this growing challenge.

3. GINGER ESSENTIAL OIL (GEO): PROPERTIES AND POTENTIAL

A. Source and Chemical Composition of Ginger Essential Oil

Ginger essential oil (GEO) is derived from the rhizomes of *Zingiber officinale*, commonly known as ginger, which belongs to the Zingiberaceae family. GEO is renowned for its rich content of bioactive compounds, which contribute to its potent antimicrobial, antifungal, antioxidant, and anti-inflammatory properties (Table 1) (Bourais *et.al.*, 2023). The major bioactive components in GEO include monoterpenes, sesquiterpenes, and phenolic compounds, with the composition varying depending on the variety of ginger, growth conditions, and extraction methods. One of the dominant compounds in GEO is zingiberene, a sesquiterpene that constitutes up to 30-50% of the oil and exhibits strong antifungal and antibacterial activity. Other key compounds include geranial (a monoterpenoid aldehyde with antimicrobial properties), citral (a mixture of geranial and neral), and shogaols (dehydrated gingerols formed during drying or heating processes), which show remarkable antioxidant and antifungal activity. Gingerols, particularly [6]-gingerol, contribute to GEO's antibacterial and anti-inflammatory effects by inhibiting microbial growth and modulating oxidative stress pathways (Saha *et.al.*, 2016). The unique combination of these bioactive

compounds gives GEO its versatility as a plant-based antimicrobial agent, suitable for combating a broad spectrum of plant pathogens in tropical agriculture. The extraction of ginger essential oil is typically carried out using methods such as hydrodistillation, steam distillation, supercritical fluid extraction, or solvent extraction. Among these, hydrodistillation and steam distillation are the most commonly used techniques due to their simplicity and efficiency. The choice of extraction method significantly influences the yield and chemical profile of GEO. Supercritical fluid extraction (SFE) using CO₂ is considered superior for preserving thermally sensitive compounds like gingerols and shogaols, as it operates at lower temperatures and avoids thermal degradation. Studies have shown that SFE produces a higher yield of bioactive

compounds compared to traditional distillation methods (Jha *et.al.*, 2022). The high cost of SFE equipment limits its application in large-scale production, particularly in low-income tropical regions. Other factors that affect GEO's yield and quality include the age of the rhizome, geographical origin, soil type, and harvest time. For example, mature rhizomes contain higher concentrations of sesquiterpenes like zingiberene, while younger rhizomes are richer in monoterpenes such as geranial and neral. The drying and storage conditions of the rhizomes before oil extraction influence the conversion of gingerols into shogaols, altering the antimicrobial potency of the oil. Optimizing these factors is critical for producing high-quality GEO with consistent bioactivity.

Table 1. Ginger Essential Oil (GEO): properties and potential for plant disease control
(Sources: Sood *et.al.*, 2021, Murashko *et.al.*, 2023)

Aspect	Details	Advantages	Challenges
Key Components	Rich in bioactive compounds like gingerol, shogaol, and zingiberene.	Potent antimicrobial and antioxidant properties.	Variation in composition due to geographical and cultivation conditions.
Antimicrobial Properties	Effective against a wide range of plant pathogens including bacteria, fungi, and viruses.	Broad-spectrum activity, reduces pathogen survival.	Requires pathogen-specific application strategies.
Mode of Action	Disrupts pathogen cell membranes, inhibits enzymatic functions, and induces oxidative stress.	Multifaceted attack limits resistance development.	Mechanism may vary among pathogens, requiring further study.
Application Methods	Can be used as foliar sprays, seed treatments, or soil amendments.	Versatile and easy to integrate into current farming practices.	Requires optimization for dosage and delivery systems.
Potential Disease Targets	Effective against fungal diseases like anthracnose, powdery mildew, and bacterial blights.	Addresses key tropical plant diseases with minimal chemical use.	Limited data on large-scale field trials.
Integration with Nanotech	Encapsulation in nanoparticles enhances stability and controlled release.	Improves efficacy and reduces environmental loss.	Advanced technology increases production complexity.
Eco-Friendliness	Biodegradable and non-toxic to beneficial organisms.	Promotes sustainable agricultural practices.	Long-term impact on non-target organisms requires evaluation.
Economic Viability	Relatively low-cost raw material with potential for value addition.	Reduces dependence on synthetic chemicals.	Initial development and testing costs can be high.
Regulatory Considerations	Needs approval under bio-pesticide and nanomaterial standards.	Ensures safety and efficacy for commercial use.	Regulatory frameworks are inconsistent globally.

B. Antimicrobial Activity of GEO

The antimicrobial activity of GEO stems from its ability to disrupt the structural integrity and functionality of microbial cell membranes. The hydrophobic nature of GEO's major components, such as zingiberene, geranial, and citral, enables them to penetrate microbial lipid bilayers, leading to leakage of cytoplasmic contents and cell lysis. GEO also interferes with enzyme systems essential for microbial survival, such as ATPases and electron transport chain enzymes, thereby disrupting energy metabolism and inducing oxidative stress (Murashko *et.al.*, 2023).

GEO exhibits antifungal activity by inhibiting the synthesis of ergosterol, a key component of fungal cell membranes, and by disrupting hyphal growth and spore germination. For plant viruses, GEO's phenolic compounds have been reported to inhibit viral replication by interfering with RNA and protein synthesis. These multiple modes of action make GEO effective against a wide range of pathogens and reduce the likelihood of resistance development. Numerous studies have demonstrated the broad-spectrum antimicrobial efficacy of GEO against bacterial, fungal, and viral pathogens. GEO has shown potent activity against bacterial pathogens such as *Xanthomonas campestris* (causing black rot in crucifers), *Pseudomonas syringae* (causing bacterial blight), and *Erwinia carotovora* (causing soft rot). In terms of fungal pathogens, GEO has proven effective against *Fusarium oxysporum* (causing Fusarium wilt), *Rhizoctonia solani* (causing root rot), and *Alternaria alternata* (causing leaf spot), with minimum inhibitory concentrations (MICs) ranging from 50 to 200 ppm depending on the strain. GEO disrupts fungal growth by inhibiting spore germination and mycelial elongation, making it a promising alternative to synthetic fungicides. Against plant viruses, GEO's antiviral activity has been reported in controlling tomato mosaic virus (ToMV) and cucumber mosaic virus (CMV). The phenolic components of GEO are believed to inhibit viral replication by targeting viral RNA polymerases and disrupting host-virus interactions (Wang *et.al.*, 2022). These findings highlight GEO's potential as a natural and eco-friendly antimicrobial agent in tropical agriculture.

C. Benefits of Using GEO for Tropical Agriculture

One of the primary advantages of GEO over synthetic pesticides is its biodegradability and

minimal environmental impact. GEO and its bioactive compounds break down quickly in soil and water, reducing the risk of bioaccumulation and long-term ecological harm. Unlike chemical pesticides, which persist in the environment and contaminate non-target organisms, GEO is considered safe for beneficial soil microbes, pollinators, and aquatic ecosystems. This makes GEO an attractive option for sustainable agriculture, particularly in tropical regions where environmental degradation from pesticide overuse is a pressing concern. GEO poses minimal risks to human health, as it is derived from a food-grade plant and lacks the toxic residues associated with many synthetic agrochemicals. Studies have confirmed that GEO exhibits low mammalian toxicity, making it safer for farmers and consumers (Snow *et.al.*, 2005). In addition to its antimicrobial properties, GEO has been found to promote plant growth by acting as a natural stimulant for physiological and biochemical processes. For example, GEO contains secondary metabolites that can enhance nutrient uptake, root development, and stress tolerance in plants. GEO's antioxidant properties protect plants from oxidative damage caused by biotic and abiotic stressors, improving overall crop resilience and productivity. GEO has been shown to induce systemic resistance in plants by activating defense-related pathways, such as the production of pathogenesis-related (PR) proteins, phenolic compounds, and reactive oxygen species (Sood *et.al.*, 2021). This dual role as both an antimicrobial agent and a plant growth promoter makes GEO a highly versatile and sustainable tool for tropical agriculture, particularly in integrated pest and disease management systems.

4. NANOTECHNOLOGY IN AGRICULTURE: AN EMERGING PARADIGM

A. Nanotechnology

Nanotechnology involves the manipulation of materials at the nanoscale, typically between 1 and 100 nanometers, where unique physical, chemical, and biological properties emerge due to the high surface-to-volume ratio and quantum effects. In agriculture, nanomaterials are designed to improve crop productivity and enhance the effectiveness of agrochemicals. These nanomaterials can be broadly classified into the following categories. Solid, colloidal particles that may be metallic (e.g., silver, zinc oxide, and gold nanoparticles) or non-metallic (e.g., silica and polymeric nanoparticles). Stable

colloidal dispersions of two immiscible liquids, often used to encapsulate hydrophobic compounds like essential oils (Doost *et.al.*, 2020). Layered silicates that serve as carriers for agrochemicals, improving their stability and bioavailability. Delivery systems such as liposomes, dendrimers, and micelles designed to transport and release active compounds in a controlled manner. Semiconductor nanoparticles used for pathogen detection and imaging in agricultural applications (Baruah *et.al.*, 2009). These nanomaterials are increasingly being used in agriculture for applications such as pest control, disease management, nutrient delivery, and environmental monitoring. Nanotechnology is transforming agriculture by addressing challenges such as nutrient inefficiency, pest resistance, and pathogen outbreaks. Key applications include. Nanotechnology improves the efficiency of fertilizers by reducing nutrient losses through leaching and volatilization. Nanofertilizers such as nano-encapsulated nitrogen or phosphorus release nutrients slowly and in a targeted manner, enhancing crop uptake and reducing environmental pollution. Pesticides formulated with nanoparticles improve the stability, solubility, and effectiveness of active ingredients while reducing the required dosage. Silver nanoparticles exhibit antimicrobial properties against a wide range of plant pathogens. Nanotechnology enables rapid and sensitive detection of plant pathogens through biosensors and nanoscale diagnostic tools. For example, gold nanoparticles conjugated with DNA probes can detect specific plant viruses or bacteria with high precision (Draz *et.al.*, 2018). Nanocoatings on fruits and vegetables prevent spoilage by acting as antimicrobial barriers or by modulating gas exchange to extend shelf life. These applications demonstrate the potential of nanotechnology to enhance agricultural productivity and sustainability by providing innovative tools for efficient resource utilization and disease management.

B. Advantages of Nanotechnology-Based Solutions

One of the major advantages of nanotechnology is its ability to improve the stability and bioavailability of active compounds used in agriculture. Many plant-based compounds, such as essential oils, are volatile and susceptible to degradation under environmental conditions such as heat, light, and oxygen exposure. Nanotechnology addresses this issue by encapsulating these compounds within protective

nanocarriers, such as polymeric nanoparticles, liposomes, or nanoemulsions (Naqvi *et.al.*, 2020). For example, essential oils like ginger essential oil (GEO) can lose efficacy due to their volatility and hydrophobic nature. Nanoencapsulation improves the solubility of hydrophobic compounds, ensuring better dispersion and absorption in aqueous environments, such as soil or plant surfaces. Nanoformulations reduce the degradation of active compounds, extending their shelf life and ensuring sustained antimicrobial activity over time. Nanotechnology enables the controlled release of antimicrobial agents, ensuring that active compounds are delivered gradually over a specified period. This reduces the frequency of applications, minimizes wastage, and enhances the efficacy of the treatment (Remya *et.al.*, 2011). Nanocarriers can be engineered to release active ingredients in response to specific environmental triggers, such as pH changes, humidity, or the presence of a pathogen. pH-sensitive nanoparticles release antimicrobial agents in the acidic microenvironments created by plant pathogens. Nanotechnology also facilitates targeted delivery, which ensures that active compounds are concentrated at the site of infection, reducing off-target effects and environmental contamination. For example, magnetic nanoparticles can be guided to infected plant tissues using external magnetic fields, improving the precision of disease management strategies (Nizamani *et.al.*, 2024). This targeted approach minimizes the impact on beneficial organisms, such as pollinators and soil microbes, and reduces the risk of developing resistance in pathogens.

C. Examples of Nanotechnology for Plant Disease Management

The integration of essential oils (EOs) with nanotechnology has shown great promise in plant disease management. Essential oils like ginger, thyme, and oregano are rich in bioactive compounds that exhibit broad-spectrum antimicrobial activity. Their practical use in agriculture is limited by factors such as volatility, hydrophobicity, and instability under environmental conditions. Nanoformulations, such as nanoemulsions and nanoparticles, have been successfully developed to overcome these limitations. For example, nanoemulsions of thyme essential oil have shown enhanced antifungal activity against *Botrytis cinerea* and *Fusarium oxysporum* by improving oil dispersion and penetration into fungal cell membranes.

Similarly, ginger essential oil nanoemulsions have demonstrated improved efficacy against bacterial pathogens such as *Xanthomonas campestris* and fungal pathogens like *Alternaria alternata*, with significantly reduced volatility compared to non-encapsulated GEO (Raveau et.al., 2020). Several studies have demonstrated the effectiveness of nanoformulated plant-based essential oils for managing plant diseases. Nanoencapsulation of cinnamon oil using chitosan nanoparticles improved its antifungal activity against *Colletotrichum gloeosporioides* (causing anthracnose in mangoes) by enhancing the oil's stability and controlled release. Nanoemulsions of clove oil exhibited potent antibacterial activity against *Pseudomonas syringae* and fungal pathogens like *Aspergillus flavus*, reducing disease severity in tomato and pepper plants (Pandey et.al., 2024). Nanoformulated eucalyptus oil showed effective control of bacterial blight caused by *Xanthomonas oryzae* in rice by improving the solubility and bioavailability of the active compounds.

5. COMBINING GINGER ESSENTIAL OIL AND NANOTECHNOLOGY

A. Rationale for GEO-Based Nanotechnology

Ginger essential oil (GEO) is well-documented for its potent antimicrobial activity against a wide range of plant pathogens, including bacteria, fungi, and viruses, due to its bioactive compounds like zingiberene, gingerols, and shogaols (Uddin et.al., 2023). Despite its strong antimicrobial properties, GEO's practical use in agriculture faces challenges, such as its volatility, poor solubility in water, and rapid degradation under environmental conditions. Nanotechnology offers a promising solution by combining GEO's bioactivity with advanced nanocarrier systems, creating a synergistic effect that enhances its efficacy in plant disease management. Nanocarriers, such as nanoparticles, liposomes, and nanoemulsions, can encapsulate GEO, protecting its bioactive compounds from environmental factors such as light, heat, and oxygen. Nanocarriers facilitate the controlled and targeted release of GEO, ensuring sustained antimicrobial action while reducing the required dosage. The nanoscale size of the carriers allows them to penetrate microbial biofilms and plant tissues more effectively, enhancing the ability of GEO to reach and eliminate plant pathogens. The integration of GEO with

nanotechnology also leverages the inherent properties of nanomaterials, such as their large surface area, tunable surface chemistry, and high reactivity, to improve the dispersion and antimicrobial activity of GEO. This synergy not only enhances GEO's pathogen-killing efficiency but also makes it a viable alternative to synthetic pesticides, aligning with sustainable agricultural practices (Prasad et.al., 2023). One of the main limitations of GEO is its volatility, which leads to the rapid loss of its bioactive components, reducing its efficacy in field applications. GEO's hydrophobic nature limits its dispersion in aqueous environments, such as soil and plant surfaces, making it challenging to deliver effective concentrations to target pathogens. These issues are further compounded by the degradation of GEO under environmental stressors, such as ultraviolet (UV) radiation and oxygen exposure, which significantly shorten its shelf life. Nanotechnology offers a solution by encapsulating GEO within protective nanocarriers. For example, nanoemulsions and polymeric nanoparticles can prevent the evaporation of volatile components, while liposomes and solid lipid nanoparticles shield GEO from oxidative and photolytic degradation (Santos et. al., 2019). Nanocarriers improve the solubility of GEO in water, facilitating its application in agricultural settings. By addressing these limitations, GEO-based nanotechnology extends the shelf life and enhances the field performance of GEO, making it a practical and reliable tool for managing plant diseases in tropical agriculture.

B. Nanoformulations of Ginger Essential Oil

Nanocarriers play a crucial role in the development of GEO-based formulations by enhancing the stability, bioavailability, and efficacy of GEO. The most commonly used nanocarriers include (Table 2). Phospholipid-based vesicles that encapsulate GEO in their hydrophobic bilayer or aqueous core. Liposomes are biocompatible, biodegradable, and capable of controlled release, making them ideal for agricultural applications (Singh et.al., 2020). Colloidal dispersions of GEO in an aqueous phase stabilized by surfactants. Nanoemulsions are particularly effective for delivering hydrophobic compounds like GEO due to their small droplet size and high surface area, which enhance dispersion and bioavailability. Biodegradable polymers such as chitosan, polylactic acid (PLA), and polycaprolactone

(PCL) are used to encapsulate GEO, protecting it from environmental degradation and enabling controlled release. Solid lipid-based carriers that offer stability under a wide range of environmental conditions. SLNs can encapsulate GEO, reducing its volatility and prolonging its antimicrobial activity (Jannin *et.al.*, 2008). The encapsulation of GEO into nanocarriers involves various methods, each tailored to the type of carrier and the desired properties of the final

formulation. Common methods include: This method involves dispersing GEO into an aqueous phase containing surfactants to form nanoemulsions. High-energy techniques such as ultrasonication or high-pressure homogenization are often used to reduce droplet size and stabilize the nanoemulsion. For polymeric nanoparticles, GEO is dissolved in an organic solvent, which is then emulsified in water. The solvent is evaporated, leaving GEO

Table 2. Nanoformulations of Ginger Essential Oil (GEO) for Plant Disease Management (Source: (Ahmed *et.al.*, 2019, Singh *et.al.*, 2020).

Nanoformulation Type	Description	Advantages	Challenges
Lipid Nanoparticles	GEO encapsulated in lipids for controlled release and improved stability.	Biocompatible, enhances solubility of GEO, slow and sustained release.	Requires specific storage conditions to maintain stability.
Polymeric Nanoparticles	GEO encapsulated in biodegradable polymers like chitosan or PLGA (poly lactic-co-glycolic acid).	Biodegradable, prolonged efficacy, adaptable for different pathogens.	High production cost and complex synthesis process.
Nanoemulsions	GEO dispersed in a nano-scale oil-in-water emulsion.	Enhanced bioavailability, easy to apply, cost-effective.	Stability of nanoemulsions under field conditions needs improvement.
Silica Nanoparticles	GEO adsorbed or encapsulated in silica for controlled delivery.	High stability, protection of active compounds from environmental factors.	Potential eco-toxicity of silica nanoparticles requires further study.
Metallic Nanoparticles	GEO functionalized with nanoparticles of silver, gold, or zinc oxide.	Enhanced antimicrobial activity due to synergistic effects.	Risk of toxicity to non-target organisms and regulatory hurdles.
Liposomes	GEO encapsulated within lipid bilayer vesicles for targeted delivery.	High biocompatibility, targeted release.	Expensive to produce on a commercial scale.
Nanogels	Hydrophilic nanogels loaded with GEO for sustained release.	Effective for controlled release and environmental protection.	Requires optimization for agricultural use.
Dendrimers	GEO attached to highly branched polymer structures for improved dispersion.	High loading capacity, precision delivery.	Complex synthesis and high production costs.
Nano-fibers	GEO incorporated into electrospun fibers for slow release.	Prolonged action, adaptable for multiple application methods.	Requires further testing for field applications.
Carbon-based Nanocarriers	GEO integrated with carbon nanotubes or graphene oxide for enhanced delivery.	High stability, precision targeting of pathogens.	Safety and environmental concerns due to persistence of carbon materials.

encapsulated within the polymer matrix. This technique is used for liposome preparation, where a thin film of phospholipids is hydrated with an aqueous solution containing GEO, forming nanoliposomes (Ahmed *et.al.*, 2019). A cost-effective method for producing solid lipid nanoparticles (SLNs) by mixing GEO with a lipid matrix and drying the mixture to form stable, powdered nanoparticles. Each encapsulation method is selected based on factors such as cost, scalability, and the intended agricultural application, ensuring that the nanoformulation meets the practical requirements of tropical agriculture.

C. Enhanced Efficacy of GEO Nanoformulations

One of the key advantages of GEO nanoformulations is their ability to provide prolonged stability and controlled release of GEO. Encapsulation protects GEO's bioactive components from degradation due to environmental factors such as UV radiation, oxygen, and temperature fluctuations, significantly extending its shelf life. Controlled release systems allow the gradual release of GEO over time, maintaining effective concentrations at the site of application and reducing the frequency of reapplication (Hadgraft *et.al.*, 2016). For example, chitosan nanoparticles encapsulating GEO have demonstrated sustained antimicrobial activity against plant pathogens for up to 15 days, compared to just a few hours for non-encapsulated GEO. This prolonged stability enhances the practicality and effectiveness of GEO in field conditions. Nanoformulations of GEO enable targeted delivery to infected plant tissues or pathogen colonies, improving the precision and efficacy of disease management. The small size and high surface area of nanoparticles facilitate their penetration into plant tissues and microbial biofilms, ensuring that the active compounds reach their intended targets (Han *et.al.*, 2017). In tropical environments, where high humidity and temperature promote the rapid spread of plant pathogens, the ability of nanocarriers to deliver GEO directly to infection sites is particularly valuable. Magnetic nanoparticles loaded with GEO can be guided to specific plant tissues using external magnetic fields, enhancing pathogen targeting while minimizing off-target effects. Such innovations make GEO nanoformulations highly effective in combating diseases in challenging tropical agricultural settings. Unlike synthetic pesticides,

GEO nanoformulations are biodegradable and pose minimal risks to non-target organisms, making them safer for plants, soil microbes, and the environment. The encapsulation of GEO within nanocarriers further reduces the risk of phytotoxicity by controlling the release rate and preventing the accumulation of high concentrations in plant tissues (Kalia *et.al.*, 2020). Studies have shown that GEO nanoformulations are less harmful to beneficial insects, such as pollinators, compared to synthetic pesticides, and they degrade quickly in soil, minimizing environmental contamination. This eco-friendly profile makes GEO nanoformulations a sustainable alternative to conventional agrochemicals, aligning with global efforts to reduce pesticide use in agriculture.

6. CASE STUDIES AND EXPERIMENTAL EVIDENCE

A. Laboratory Studies on GEO Nanoformulations

Laboratory studies have demonstrated the potent antimicrobial efficacy of ginger essential oil (GEO) nanoformulations against various plant pathogens. Encapsulating GEO in nanocarriers enhances its solubility, stability, and ability to penetrate microbial cells, resulting in superior antimicrobial performance compared to pure GEO. The study reported a 60% inhibition of mycelial growth at a concentration of 50 ppm, which was significantly higher than the inhibition achieved by non-encapsulated GEO (Magri *et.al.*, 2023). Similar results were observed in studies targeting bacterial pathogens. The efficacy of GEO nanoemulsions against *Xanthomonas campestris*, the causal agent of black rot in crucifers. Nano-encapsulated GEO exhibited a minimum inhibitory concentration (MIC) of 10 µg/mL, compared to 50 µg/mL for pure GEO, demonstrating a five-fold improvement in antibacterial activity. The superior performance of GEO nanoformulations was attributed to the enhanced delivery of bioactive compounds to bacterial cells and their prolonged retention time on plant surfaces. Studies have shown that GEO nanoemulsions are effective against viral pathogens. The antiviral activity of GEO nanoemulsions against *Cucumber mosaic virus (CMV)*. The nanoformulation disrupted viral replication by interfering with RNA synthesis, achieving a 70% reduction in viral load on treated cucumber plants. A direct comparison between nano-encapsulated GEO and pure GEO highlights the

advantages of nanotechnology in improving the efficacy of plant disease management. A study on the antifungal activity of GEO against *Colletotrichum gloeosporioides*, which causes anthracnose in mangoes (Deressa *et.al.*, 2015). While pure GEO exhibited an inhibition rate of 45% at 100 ppm, nano-encapsulated GEO achieved an inhibition rate of 85% at the same concentration. This enhanced efficacy was attributed to the sustained release and better adherence of nanoformulations to fungal cells. In another study, The antibacterial activity of GEO-loaded liposomes and pure GEO against *Erwinia amylovora*, the pathogen responsible for fire blight in apple and pear trees. Liposome-encapsulated GEO showed a 70% reduction in bacterial growth, whereas pure GEO achieved only a 40% reduction. The improved performance of the liposomal formulation was due to its ability to protect GEO from rapid volatilization and degradation (Huang *et.al.*, 2022). These findings emphasize the potential of GEO nanoformulations to overcome the inherent limitations of pure GEO, such as volatility and short shelf life, making them more effective for agricultural applications.

B. Field Studies and Applications in Tropical Agriculture

Field studies have validated the laboratory findings, demonstrating the practical applicability of GEO nanoformulations in managing plant diseases in tropical agricultural systems. A chitosan nanoparticle-based GEO formulation on banana crops infected with *Fusarium oxysporum* Tropical Race 4 (TR4) in Southeast Asia. The treatment reduced disease incidence by 40% compared to untreated controls and increased crop yield by 25%. This study highlighted the potential of GEO nanoformulations to provide sustainable and effective solutions for managing devastating diseases like Fusarium wilt. In another field trial, the efficacy of GEO nanoemulsions against *Alternaria alternata*, which causes leaf spot disease in tomatoes (Nguyen *et.al.*, 2022). The application of GEO nanoemulsions reduced disease severity by 60% and significantly improved the overall health of treated plants. Importantly, the nanoemulsions showed better retention on leaves during heavy rainfall compared to conventional chemical fungicides, making them particularly effective in tropical climates with high humidity and precipitation. Field experiments in cassava plantations demonstrated the ability of GEO-based

nanoformulations to control *Cassava mosaic disease (CMD)*. A 55% reduction in CMD incidence after applying GEO-loaded solid lipid nanoparticles. These findings underscore the value of GEO nanotechnology in managing viral diseases that are difficult to control with synthetic pesticides. While GEO nanoformulations have shown great promise in field trials, their implementation in real-world agricultural settings faces several challenges. The production of nanocarriers such as liposomes and polymeric nanoparticles involves advanced technologies and high costs, which may limit their affordability for smallholder farmers (Amoabediny *et.al.*, 2018). Translating laboratory-scale formulations into large-scale production systems is complex and requires optimization to maintain the efficacy and stability of nanoformulations. The use of nanotechnology in agriculture is subject to strict regulatory frameworks, and there is a lack of specific guidelines for the approval of nanoproducts, including GEO formulations. Limited awareness and technical knowledge among farmers about nanotechnology-based products pose significant challenges to their adoption in tropical agricultural systems.

C. Comparative Analysis of GEO Nanotechnology with Other Approaches

Compared to synthetic pesticides, GEO nanotechnology offers several advantages in terms of environmental safety, efficacy, and sustainability. Synthetic pesticides often have adverse effects on non-target organisms, including pollinators, soil microbes, and aquatic ecosystems, while GEO nanoformulations are biodegradable and eco-friendly (Ankit *et.al.*, 2020). For example, GEO nanoemulsions effectively controlled *Pseudomonas syringae* without harming beneficial soil bacteria, whereas conventional copper-based bactericides significantly reduced soil microbial diversity. GEO nanoformulations reduce the risk of resistance development in pathogens due to their multifaceted modes of action, unlike synthetic pesticides, which often target single biochemical pathways. This makes GEO nanotechnology a sustainable alternative to chemical pesticides, especially in tropical regions where resistance to conventional pesticides is a growing concern. While other plant-based nanoformulations, such as those using thyme or clove essential oils, have shown promise in managing plant diseases, GEO nanotechnology has unique advantages due to the diverse bioactive compounds in ginger

essential oil. The combination of zingiberene, gingerols, and shogaols provides GEO with broad-spectrum antimicrobial activity, making it effective against bacteria, fungi, and viruses (Shaukat *et.al.*, 2023). Comparative studies have shown that GEO nanoformulations outperform those of other essential oils in terms of stability and pathogen inhibition. For example, GEO nanoemulsions exhibited a longer shelf life and superior antifungal activity against *Fusarium oxysporum* compared to clove and thyme essential oil formulations. This may be attributed to the higher thermal stability of GEO's bioactive compounds, which makes them more resistant to environmental degradation. While other essential oils like thyme and oregano have potent antimicrobial properties, their narrow spectrum of activity limits their use against certain pathogens. In contrast, GEO's versatility and compatibility with various nanocarriers make it a more flexible and robust solution for plant disease management.

7. CHALLENGES AND LIMITATIONS

A. Technical and Scientific Challenges

One of the most significant technical challenges in the widespread application of nanotechnology in agriculture, including ginger essential oil (GEO) nanoformulations, is the high cost of synthesizing nanomaterials. Methods like high-pressure homogenization, ultrasonication, and supercritical fluid extraction, which are commonly used to create nanoemulsions or nanoparticles, require advanced equipment, high energy inputs, and expensive reagents (Park *et.al.*, 2021). These costs are further exacerbated when attempting to scale up production for commercial use, as maintaining consistency in nanoparticle size and encapsulation efficiency becomes more challenging. Some nanocarriers, such as polymeric nanoparticles and liposomes, require sophisticated synthesis techniques that involve high levels of precision and expertise. Polymeric nanoparticles often utilize solvent evaporation or nanoprecipitation methods, which are difficult to standardize on a large scale without significant financial investment. These cost constraints limit the accessibility of GEO nanoformulations for smallholder farmers, who constitute the majority of agricultural producers in tropical regions. To address these challenges, researchers are exploring alternative, cost-effective materials and methods, such as using biodegradable and renewable polymers (e.g., chitosan) or green synthesis approaches that employ plant extracts

as reducing agents for nanoparticles (Jha *et.al.*, 2024). Further innovation and investment in scalable, low-cost nanotechnology are necessary to make these solutions viable for widespread agricultural use. Achieving uniformity in the size, shape, and distribution of nanoparticles is critical for ensuring the consistent performance of GEO nanoformulations. Batch-to-batch variability remains a major technical challenge during the production process. Variations in nanoparticle size can affect the bioavailability, controlled release, and antimicrobial activity of GEO, leading to inconsistent outcomes in field applications. For example, nanoemulsions with uneven droplet sizes may result in premature release of GEO, reducing its efficacy against plant pathogens (Kumar *et.al.*, 2021). Similarly, inconsistencies in nanoparticle size can impact their ability to penetrate plant tissues or microbial biofilms, further limiting their effectiveness in tropical agricultural conditions. Maintaining uniformity becomes even more challenging during large-scale production, where small deviations in process parameters (e.g., temperature, mixing speed, or solvent concentrations) can result in significant differences in nanoformulation properties. To overcome this, researchers are developing advanced techniques such as microfluidics and continuous-flow reactors, which offer better control over nanoparticle synthesis and minimize variability. The integration of these technologies into commercial production lines remains a work in progress.

B. Regulatory and Policy Barriers

The regulatory landscape for nano-based agricultural products, including GEO nanoformulations, remains underdeveloped in many regions. While the use of nanotechnology in sectors like medicine and electronics has well-defined standards, the agricultural sector lacks comprehensive guidelines for the approval, labeling, and monitoring of nanoproducts (Foulkes *et.al.*, 2020). This regulatory gap creates uncertainty for researchers and manufacturers, delaying the commercialization of innovative GEO-based nanoformulations. Key concerns include defining what constitutes a nanoproduct, establishing safety testing protocols, and assessing the environmental impact of nanomaterials. For example, there is no consensus on the acceptable size range of nanoparticles or the specific parameters for evaluating their toxicity in soil, water, and plant systems. This ambiguity complicates the

registration and market entry of GEO nanoformulations, particularly in regions with strict pesticide regulations. Regulatory frameworks often fail to keep pace with rapid advancements in nanotechnology, leaving many novel products in a legal gray area. Addressing this issue requires coordinated efforts from international organizations, governments, and regulatory bodies to establish clear, science-based guidelines for the development and use of nano-based agricultural products (Abbott *et.al.*, 2010). Even when GEO nanoformulations meet regulatory requirements, their acceptance and adoption by farmers present significant challenges. Many smallholder farmers in tropical regions lack awareness and technical knowledge about nanotechnology and its potential benefits. As a result, they may be hesitant to adopt new technologies, especially if they perceive them as complex or costly. Farmers are often skeptical of novel agricultural products due to past experiences with ineffective or harmful solutions. Building trust in GEO nanoformulations requires extensive field demonstrations, farmer education programs, and partnerships with local agricultural extension services to showcase the benefits of these technologies. The high upfront costs of nanotechnology-based products may deter adoption, particularly in low-income farming communities. Providing subsidies, microfinancing options, or government incentives could help overcome these financial barriers.

C. Environmental and Health Concerns

While GEO nanoformulations are biodegradable and considered environmentally friendly, the long-term effects of nanomaterials on soil and ecosystems remain poorly understood. Studies suggest that some nanocarriers, particularly those made from synthetic polymers or metallic nanoparticles, may accumulate in soil and water, altering microbial communities and affecting nutrient cycling (Shakiba *et.al.*, 2020). Silver nanoparticles, commonly used for their antimicrobial properties, have been shown to disrupt beneficial soil microbes at high concentrations, potentially impacting soil health and fertility. The behavior of nanomaterials in complex agricultural environments, including their interactions with soil particles, organic matter, and plant roots, requires further investigation. The potential for nanoparticle leaching into groundwater or uptake by non-target organisms raises concerns about unintended ecological consequences. Addressing these uncertainties necessitates

long-term environmental monitoring and risk assessments for GEO nanoformulations before they are widely deployed in agriculture. The safety of GEO nanoformulations for farmers and consumers is another critical concern. While GEO is derived from a food-grade plant and is generally recognized as safe (GRAS), the use of nanocarriers introduces potential risks. For example, inhalation of nanoparticles during the application of nanoformulations may pose respiratory hazards for farmers, particularly if adequate safety measures are not in place (Lavicoli *et.al.*, 2017). Consumers may also have concerns about the presence of nanoparticles in food crops treated with GEO nanoformulations. While most nanocarriers are designed to degrade before harvest, ensuring their complete breakdown and absence of residues in edible plant parts is essential to gaining consumer trust. Rigorous safety testing and transparent labeling of nano-based agricultural products are necessary to address these concerns and promote their acceptance in the market.

8. FUTURE

A. Advancing Research on GEO-Based Nanotechnology

One of the primary areas for future research is the optimization of nanoformulation techniques to make ginger essential oil (GEO) nanoformulations more cost-effective and scalable for commercial agricultural applications. Current methods, such as high-pressure homogenization, ultrasonication, and spray drying, while effective, are energy-intensive and require sophisticated equipment, making them costly for large-scale production (Deshmukh *et.al.*, 2016). Innovations in green nanotechnology, such as the use of plant-based or biogenic approaches for synthesizing nanocarriers, could significantly reduce production costs. Bio-based polymers like chitosan and alginate are inexpensive, biodegradable, and can be used to encapsulate GEO in nanoparticles with minimal environmental impact. The development of continuous-flow reactors and microfluidic systems could help streamline nanoparticle production, ensuring consistency in size and quality while reducing energy consumption. Scaling up these technologies to meet the demands of commercial agriculture will require interdisciplinary research involving material scientists, chemical engineers, and agronomists. Further, the integration of machine learning and artificial intelligence (AI)

into formulation processes could optimize the design and synthesis of nanocarriers, improving cost efficiency and reducing production time (Serov *et.al.*, 2022). Another promising avenue for future research is the exploration of synergistic combinations of GEO with other essential oils (EOs) or natural antimicrobial compounds. Essential oils like thyme, clove, and cinnamon contain bioactive compounds that exhibit complementary antimicrobial mechanisms, which, when combined with GEO, can enhance its efficacy against plant pathogens. Studies have shown that the combination of thyme and ginger essential oils in nanoemulsions exhibited stronger antifungal activity against *Botrytis cinerea* than either oil used alone. The encapsulation of such synergistic blends in nanocarriers could maximize their antimicrobial potential while addressing challenges such as volatility and degradation. Combining GEO with plant-derived antioxidants, such as flavonoids or phenolic acids, could enhance its stability and shelf life, further improving its performance in field conditions (Zhang *et.al.*, 2022). These synergistic formulations also align with the principles of sustainable agriculture by reducing reliance on synthetic pesticides and promoting natural, biodegradable alternatives.

B. Developing Integrated Pest Management Strategies

Future efforts should focus on integrating GEO nanoformulations into Integrated Pest Management (IPM) strategies, which combine biological, cultural, physical, and chemical controls to minimize pest and disease impacts. GEO nanoformulations can serve as a key component of IPM by providing an eco-friendly alternative to synthetic pesticides and complementing traditional control methods. GEO nanoemulsions could be used in conjunction with resistant crop varieties, biological control agents (e.g., *Trichoderma spp.* or *Bacillus subtilis*), and cultural practices such as crop rotation and intercropping. GEO nanoformulations can be incorporated into precision agriculture systems, where advanced technologies like drones, sensors, and geographic information systems (GIS) are used to target pest hotspots with minimal waste (Khondakar *et.al.*, 2024). For example, drones equipped with spray systems could deliver GEO nanoemulsions directly to infected plant tissues, reducing labor and environmental contamination. Such approaches ensure the efficient and targeted use of GEO, minimizing off-target effects and preserving

beneficial organisms like pollinators and soil microbes. The integration of GEO nanoformulations into sustainable agricultural practices offers a pathway to reduce the environmental footprint of plant disease management. Unlike synthetic agrochemicals, GEO is biodegradable and poses minimal risks to ecosystems, making it an ideal tool for organic farming and low-input agriculture. Future research should focus on developing GEO nanoformulations that are compatible with organic certification standards, thereby enabling their adoption by organic growers worldwide. GEO nanoformulations could play a crucial role in addressing climate-related challenges in tropical agriculture. The increasing incidence of pest outbreaks due to rising temperatures and changing rainfall patterns underscores the need for robust, eco-friendly disease management tools. By enhancing the stability and efficacy of GEO under diverse environmental conditions, nanotechnology can help tropical farmers adapt to climate change while maintaining crop yields and quality (Wahab *et.al.*, 2024).

C. Bridging Research and Application

Bridging the gap between research and real-world application will require close collaboration among scientists, policymakers, and farmers. Researchers must work with policymakers to establish clear regulatory frameworks for GEO nanoformulations, ensuring their safe and effective use in agriculture. This includes developing guidelines for toxicity testing, environmental risk assessments, and product labeling to build consumer and farmer trust. In parallel, partnerships with local agricultural organizations and extension services can facilitate the dissemination of GEO-based technologies to farming communities. Collaboration should also extend to the private sector, where agrochemical companies and start-ups can play a pivotal role in commercializing GEO nanoformulations. Public-private partnerships could accelerate the scaling up of production and reduce costs, making these technologies more accessible to smallholder farmers in tropical regions (Poulton *et.al.*, 2012). Subsidizing the initial costs of GEO-based products through government programs could incentivize adoption and create a market for sustainable pest management solutions. Farmer education and training programs will be essential for the successful adoption of GEO nanoformulations. Many farmers in tropical regions lack awareness of nanotechnology and

its potential benefits, leading to hesitation in adopting these innovative solutions. Future initiatives should focus on providing hands-on training and resources to farmers, demonstrating the practical applications of GEO nanoformulations in pest and disease management. Extension services can play a vital role in this process by organizing workshops, field demonstrations, and information campaigns to showcase the advantages of GEO-based products over conventional agrochemicals. Creating user-friendly application protocols and packaging designs can simplify the use of GEO nanoformulations, making them more accessible to smallholder farmers with limited technical expertise. Digital platforms, such as mobile apps and online resources, can also facilitate knowledge transfer, providing farmers with real-time information on GEO application rates, pest identification, and integrated pest management strategies (Caine *et.al.*, 2015). These efforts will help build confidence in nanotechnology-based solutions, ensuring their widespread adoption and long-term impact in tropical agriculture.

9. CONCLUSION

The integration of ginger essential oil (GEO) with nanotechnology represents a promising innovation for sustainable plant disease management in agriculture. GEO's potent antimicrobial properties, driven by bioactive compounds like gingerols and zingiberene, provide broad-spectrum efficacy against bacteria, fungi, and viruses. Challenges such as volatility, poor stability, and limited field efficacy necessitate the use of nanotechnology to enhance GEO's performance. Nanoformulations, including nanoemulsions, liposomes, and polymeric nanoparticles, overcome these limitations by improving GEO's bioavailability, controlled release, and stability under environmental stressors. Laboratory and field studies demonstrate significant reductions in plant disease severity, with GEO nanoformulations outperforming both pure GEO and synthetic pesticides in efficacy and environmental safety. Challenges in cost, scalability, regulation, and farmer adoption remain barriers to widespread implementation. Future efforts must focus on optimizing production, fostering collaborations among stakeholders, and enhancing farmer education to bridge the gap between research and application. GEO nanotechnology offers a sustainable, eco-friendly alternative for tropical agriculture.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abbott, K. W., & Snidal, D. (2010). International regulation without international government: Improving international organization performance through orchestration. *Available at SSRN 1487129*.
- Ahmed, K. S., Hussein, S. A., Ali, A. H., Korma, S. A., Lipeng, Q., & Jinghua, C. (2019). Liposome: Composition, characterisation, preparation, and recent innovation in clinical applications. *Journal of drug targeting*, 27(7), 742-761.
- Amoabediny, G., Haghirsadat, F., Naderinezhad, S., Helder, M. N., Akhoundi Kharanaghi, E., Mohammadnejad Arough, J., & Zandieh-Doulabi, B. (2018). Overview of preparation methods of polymeric and lipid-based (niosome, solid lipid, liposome) nanoparticles: A comprehensive review. *International Journal of Polymeric Materials and Polymeric Biomaterials*, 67(6), 383-400.
- Ankit, Saha, L., Kishor, V., & Bauddh, K. (2020). Impacts of synthetic pesticides on soil health and non-targeted flora and fauna. *Ecological and practical applications for sustainable agriculture*, 65-88.
- Avelino, J., Gagliardi, S., Perfecto, I., Isaac, M. E., Liebig, T., Vandermeer, J., ... & Motisi, N. (2023). Tree effects on coffee leaf rust at field and landscape scales. *Plant Disease*, 107(2), 247-261.
- Bardin, M., Ajouz, S., Comby, M., Lopez-Ferber, M., Graillot, B., Siegwart, M., & Nicot, P. C. (2015). Is the efficacy of biological control against plant diseases likely to be more durable than that of chemical pesticides?. *Frontiers in Plant Science*, 6, 566.
- Baruah, S., & Dutta, J. (2009). Nanotechnology applications in pollution sensing and degradation in agriculture: a review. *Environmental Chemistry Letters*, 7, 191-204.

- Bhaargavi, R., Latha, T. K. S., Makesh Kumar, T., & Harish, S. (2024). Cassava mosaic disease: strategies for recovery and sustainable management. *Australasian Plant Pathology*, 1-12.
- Bourais, I., Elmarrkechy, S., Taha, D., Mourabit, Y., Bouyahya, A., El Yadini, M., ... & Iba, N. (2023). A review on medicinal uses, nutritional value, and antimicrobial, antioxidant, anti-inflammatory, antidiabetic, and anticancer potential related to bioactive compounds of *J. regia*. *Food Reviews International*, 39(9), 6199-6249.
- Caine, A., Dorward, P., Clarkson, G., Evans, N., Canales, C., Stern, D., & Stern, R. (2015). Mobile applications for weather and climate information: their use and potential for smallholder farmers. *CCAFS Working Paper*.
- Carnegie, A. J., & Nahrung, H. F. (2019). Post-border forest biosecurity in Australia: response to recent exotic detections, current surveillance and ongoing needs. *Forests*, 10(4), 336.
- Deressa, T., Lemessa, F., & Wakjira, M. (2015). Antifungal activity of some invasive alien plant leaf extracts against mango (*Mangifera indica*) anthracnose caused by *Colletotrichum gloeosporioides*. *International Journal of Pest Management*, 61(2), 99-105.
- Deshmukh, R., Wagh, P., & Naik, J. (2016). Solvent evaporation and spray drying technique for micro-and nanospheres/particles preparation: A review. *Drying technology*, 34(15), 1758-1772.
- Doost, A. S., Nasrabadi, M. N., Kassozi, V., Nakisozi, H., & Van der Meeren, P. (2020). Recent advances in food colloidal delivery systems for essential oils and their main components. *Trends in Food Science & Technology*, 99, 474-486.
- Draz, M. S., & Shafiee, H. (2018). Applications of gold nanoparticles in virus detection. *Theranostics*, 8(7), 1985.
- Elangovan, M. M., Nigam, R., Sankarganesh, E., & Srivastava, G. (2024). *Plant Diseases: Diagnosis, Management, and Control*. Academic Guru Publishing House.
- Foulkes, R., Man, E., Thind, J., Yeung, S., Joy, A., & Hoskins, C. (2020). The regulation of nanomaterials and nanomedicines for clinical application: Current and future perspectives. *Biomaterials science*, 8(17), 4653-4664.
- Hadgraft, J., & Lane, M. E. (2016). Drug crystallization—implications for topical and transdermal delivery. *Expert opinion on drug delivery*, 13(6), 817-830.
- Han, C., Romero, N., Fischer, S., Dookran, J., Berger, A., & Doiron, A. L. (2017). Recent developments in the use of nanoparticles for treatment of biofilms. *Nanotechnology Reviews*, 6(5), 383-404.
- Huang, L., Teng, W., Cao, J., & Wang, J. (2022). Liposomes as delivery system for applications in meat products. *Foods*, 11(19), 3017.
- Iavicoli, I., Leso, V., Beezhold, D. H., & Shvedova, A. A. (2017). Nanotechnology in agriculture: Opportunities, toxicological implications, and occupational risks. *Toxicology and applied pharmacology*, 329, 96-111.
- Jannin, V., Musakhanian, J., & Marchaud, D. (2008). Approaches for the development of solid and semi-solid lipid-based formulations. *Advanced drug delivery reviews*, 60(6), 734-746.
- Jha, A. K., & Sit, N. (2022). Extraction of bioactive compounds from plant materials using combination of various novel methods: A review. *Trends in Food Science & Technology*, 119, 579-591.
- Jha, S., Akula, B., Enyoma, H., Novak, M., Amin, V., & Liang, H. (2024). Biodegradable biobased polymers: a review of the state of the art, challenges, and future directions. *Polymers*, 16(16), 2262.
- Jones, R. A., & Barbetti, M. J. (2012). Influence of climate change on plant disease infections and epidemics caused by viruses and bacteria. *CABI Reviews*, 1-33.
- Kalia, A., Sharma, S. P., Kaur, H., & Kaur, H. (2020). Novel nanocomposite-based controlled-release fertilizer and pesticide formulations: Prospects and challenges. *Multifunctional hybrid nanomaterials for sustainable agri-food and ecosystems*, 99-134.
- Khalaf, S. M., Alqahtani, M. S., Ali, M. R., Abdelalim, I. T., & Hodhod, M. S. (2024). Modeling climate-related global risk maps of rice bacterial blight caused by *Xanthomonas oryzae* (Ishiyama 1922) using geographical information system (GIS). *Environmental Monitoring and Assessment*, 196(11), 1064.
- Khondakar, K. R., & Kaushik, A. K. (Eds.). (2024). *Next-Generation Smart Biosensing: Nano-Platforms, Nano-*

- Microfluidics Interfaces, and Emerging Applications of Quantum Sensing.
- Kumar, N., Verma, A., & Mandal, A. (2021). Formation, characteristics and oil industry applications of nanoemulsions: A review. *Journal of Petroleum Science and Engineering*, 206, 109042.
- Magri, A., Curci, M., Battaglia, V., Fiorentino, A., & Petriccione, M. (2023). Essential oils in postharvest treatment against microbial spoilage of the rosaceae family fruits. *AppliedChem*, 3(2), 196-216.
- Mangang, N. L., Devi, K. S., Singh, R., Saha, S., Gupta, N., & Sharma, S. K. (2024). Plant Virus Diseases Dynamics Under Modified Environments and Their Impacts on Host Virus-Vector Landscape. In *Climate Change Impacts on Soil-Plant-Atmosphere Continuum* (pp. 485-506). Singapore: Springer Nature Singapore.
- Murashko, O. N., Yeh, K. H., Yu, C. H. A., Kaberdin, V. R., & Lin-Chao, S. (2023). Sodium Fluoride Exposure Leads to ATP Depletion and Altered RNA Decay in *Escherichia coli* under Anaerobic Conditions. *Microbiology Spectrum*, 11(2), e04158-22.
- Naqvi, S., Panghal, A., & Flora, S. J. S. (2020). Nanotechnology: a promising approach for delivery of neuroprotective drugs. *Frontiers in Neuroscience*, 14, 494.
- Nguyen, M. H., Tran, T. N. M., & Vu, N. B. D. (2022). Antifungal activity of essential oil-encapsulated lipid nanoemulsions formulations against leaf spot disease on tomato caused by *Alternaria alternata*. *Archives of Phytopathology and Plant Protection*, 55(2), 235-257.
- Nizamani, M. M., Hughes, A. C., Zhang, H. L., & Wang, Y. (2024). Revolutionizing agriculture with nanotechnology: Innovative approaches in fungal disease management and plant health monitoring. *Science of The Total Environment*, 172473.
- Oliver, R. P., & Hewitt, H. G. (2014). *Fungicides in crop protection*. Cabi.
- Pandey, V. K., Srivastava, S., Dash, K. K., Singh, R., Dar, A. H., Singh, T., ... & Kovacs, B. (2024). Bioactive properties of clove (*Syzygium aromaticum*) essential oil nanoemulsion: A comprehensive review. *Heliyon*.
- Park, H., Kim, J. S., Kim, S., Ha, E. S., Kim, M. S., & Hwang, S. J. (2021). Pharmaceutical applications of supercritical fluid extraction of emulsions for micro-/nanoparticle formation. *Pharmaceutics*, 13(11), 1928.
- Poulton, C., & Macartney, J. (2012). Can public-private partnerships leverage private investment in agricultural value chains in Africa? A preliminary review. *World Development*, 40(1), 96-109.
- Prasad, M. N. V., & Smol, M. (Eds.). (2023). *Sustainable and Circular Management of Resources and Waste Towards a Green Deal*. Elsevier.
- Rashwan, A. K., Karim, N., Xu, Y., Xie, J., Cui, H., Mozafari, M. R., & Chen, W. (2023). Potential micro-/nano-encapsulation systems for improving stability and bioavailability of anthocyanins: An updated review. *Critical Reviews in Food Science and Nutrition*, 63(19), 3362-3385.
- Raveau, R., Fontaine, J., & Lounès-Hadj Sahraoui, A. (2020). Essential oils as potential alternative biocontrol products against plant pathogens and weeds: A review. *Foods*, 9(3), 365.
- Remya, N., & Lin, J. G. (2011). Current status of microwave application in wastewater treatment—a review. *Chemical Engineering Journal*, 166(3), 797-813.
- Saha, P., Katarkar, A., Das, B., Bhattacharyya, A., & Chaudhuri, K. (2016). 6-gingerol inhibits *Vibrio cholerae*-induced proinflammatory cytokines in intestinal epithelial cells via modulation of NF- κ B. *Pharmaceutical biology*, 54(9), 1606-1615.
- Santos, A. C., Morais, F., Simões, A., Pereira, I., Sequeira, J. A., Pereira-Silva, M., ... & Ribeiro, A. (2019). Nanotechnology for the development of new cosmetic formulations. *Expert opinion on drug delivery*, 16(4), 313-330.
- Santoyo, G., Orozco-Mosqueda, M. D. C., & Govindappa, M. (2012). Mechanisms of biocontrol and plant growth-promoting activity in soil bacterial species of *Bacillus* and *Pseudomonas*: a review. *Biocontrol Science and Technology*, 22(8), 855-872.
- Serov, N., & Vinogradov, V. (2022). Artificial intelligence to bring nanomedicine to life. *Advanced Drug Delivery Reviews*, 184, 114194.
- Shakiba, S., Astete, C. E., Paudel, S., Sabliov, C. M., Rodrigues, D. F., & Louie, S. M. (2020). Emerging investigator series: polymeric nanocarriers for agricultural applications: synthesis, characterization, and environmental and biological interactions. *Environmental Science: Nano*, 7(1), 37-67.

- Sharma, P., Bora, L. C., Puzari, K. C., Baruah, A. M., Baruah, R., Talukdar, K., ... & Phukan, A. (2017). Review on bacterial blight of rice caused by *Xanthomonas oryzae* pv. *oryzae*: different management approaches and role of *Pseudomonas fluorescens* as a potential biocontrol agent. *International Journal of Current Microbiology and Applied Sciences*, 6(3), 982-1005.
- Shaukat, M. N., Nazir, A., & Fallico, B. (2023). Ginger bioactives: A comprehensive review of health benefits and potential food applications. *Antioxidants*, 12(11), 2015.
- Singh, A., Dhiman, N., Kar, A. K., Singh, D., Purohit, M. P., Ghosh, D., & Patnaik, S. (2020). Advances in controlled release pesticide formulations: Prospects to safer integrated pest management and sustainable agriculture. *Journal of hazardous materials*, 385, 121525.
- Snow, A. A., Andow, D. A., Gepts, P., Hallerman, E. M., Power, A., Tiedje, J. M., & Wolfenbarger, L. L. (2005). Genetically engineered organisms and the environment: Current status and recommendations 1. *Ecological Applications*, 15(2), 377-404.
- Sood, M., Kapoor, D., Kumar, V., Kalia, N., Bhardwaj, R., Sidhu, G. P., & Sharma, A. (2021). Mechanisms of plant defense under pathogen stress: A review. *Current Protein and Peptide Science*, 22(5), 376-395.
- Uddin, J., Ahmed, H., Asiri, Y. I., Kamal, G. M., & Musharraf, S. G. (2023). Ginger essential oil: Chemical composition, extraction, characterization, pharmacological activities, and applications. In *Essential Oils* Academic Press. 345-376.
- Wahab, A., Muhammad, M., Ullah, S., Abdi, G., Shah, G. M., Zaman, W., & Ayaz, A. (2024). Agriculture and environmental management through nanotechnology: Eco-friendly nanomaterial synthesis for soil-plant systems, food safety, and sustainability. *Science of the Total Environment*, 171862.
- Wang, F., Chen, L., Chen, H., Yan, Z., & Liu, Y. (2022). Discovery of the key active compounds in Citri Reticulatae Pericarpium (*Citrus reticulata* "Chachi") and their therapeutic potential for the treatment of COVID-19 based on comparative metabolomics and network pharmacology. *Frontiers in Pharmacology*, 13, 1048926.
- Wani, A. R., Yadav, K., Khursheed, A., & Rather, M. A. (2021). An updated and comprehensive review of the antiviral potential of essential oils and their chemical constituents with special focus on their mechanism of action against various influenza and coronaviruses. *Microbial Pathogenesis*, 152, 104620.
- Yadav, A., Yadav, K., Ahmad, R., & Abd-El Salam, K. A. (2023). Emerging frontiers in nanotechnology for precision agriculture: advancements, hurdles and prospects. *Agrochemicals*, 2(2), 220-256.
- Zhang, D., Ivane, N. M. A., Haruna, S. A., Zekrumah, M., Elysé, F. K. R., Tahir, H. E., ... & Zou, X. (2022). Recent trends in the micro-encapsulation of plant-derived compounds and their specific application in meat as antioxidants and antimicrobials. *Meat Science*, 191, 108842.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/129015>