



Effect of Sowing Date and Plant Growth Regulators on Growth and Yield Attributes of Wheat (*Triticum aestivum* L.) Grown at AEZ-9 of Bangladesh

Md. Sohel Rana ^a, Masuma Ahmed ^b,
Md Roconuzzaman Nasim ^a, Md Maruf Hossain ^a,
Nilufer Yasmin Jolly ^b, Kazi Rifat Shahrin ^c,
Mohammad Amdadul Hoque ^d, Md. Abdul Kader ^b
and Ahmed Khairul Hasan ^{b*}

^a Department of Soil Science, Faculty of Agriculture, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh.

^b Department of Agronomy, Faculty of Agriculture, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh.

^c Department of Biotechnology, Faculty of Agriculture, Sher-E-Bangla Agricultural University, Dhaka-1207, Bangladesh.

^d Department of Agricultural Extension Education, Faculty of Agriculture, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajrcs/2024/v9i4312>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/126457>

Original Research Article

Received: 10/09/2024
Accepted: 12/11/2024
Published: 15/11/2024

*Corresponding author: E-mail: akhasan@bau.edu.bd;

Cite as: Rana, Md. Sohel, Masuma Ahmed, Md Roconuzzaman Nasim, Md Maruf Hossain, Nilufer Yasmin Jolly, Kazi Rifat Shahrin, Mohammad Amdadul Hoque, Md. Abdul Kader, and Ahmed Khairul Hasan. 2024. "Effect of Sowing Date and Plant Growth Regulators on Growth and Yield Attributes of Wheat (*Triticum Aestivum* L.) Grown at AEZ-9 of Bangladesh". *Asian Journal of Research in Crop Science* 9 (4):215-27. <https://doi.org/10.9734/ajrcs/2024/v9i4312>.

ABSTRACT

A field experiment was conducted at the Agronomy Field Laboratory, Bangladesh Agricultural University, Mymensingh, from November 2019 to March 2020 to assess the combined effects of sowing date and plant growth regulators (PGRs) on the growth and yield of wheat cv *BARI Gom-33*. The experiment was conducted in a split-plot design with three replications, with sowing dates in the main plots and PGRs in the sub-plots, totaling 36 plots (12 treatments × 3 replications). Results of the study revealed that sowing on 5th December with Gibberellic Acid (GA₃) application increased grain yield by 25.62% over the control. This sowing date also produced 7.59% and 12.83% higher yields than early and late sowing dates, respectively. The results suggest that sowing on 5th December combined with GA₃ application effectively optimizes wheat growth and yield, providing a promising approach for enhancing wheat production and supporting sustainable agriculture.

Keywords: Yield; BARI; Split-plot; sustainable agriculture; IAA; GA₃; NAA.

1. INTRODUCTION

Wheat (*Triticum aestivum* L.) is a cornerstone of global food security, providing essential nutrition and economic support to millions of people worldwide. Accounting for approximately 30% of the world's grain production, wheat serves as a staple food in over 40 countries, supplying vital calories and protein to a significant portion of the global population [1,2]. However, as demand for wheat continues to grow, so does the need to improve production efficiency and yield quality, especially in regions facing climate variability and resource constraints [3]. Climate change, along with fluctuating weather patterns and seasonal temperatures, introduces further complexities to wheat production [4]. Thus, adopting sustainable agronomic practices that optimize wheat growth and maximize yield is essential for sustaining and enhancing global food security.

One of the most influential factors in wheat yield optimization is managing environmental conditions, particularly through the timing of sowing. Optimal sowing timing aligns wheat's vegetative and reproductive phases with favorable environmental conditions, such as temperature, photoperiod, and rainfall, leading to improved plant height, spike length, and grain filling [5]. Indeed, research shows that optimal combinations of sowing dates and seeding rates can increase yields by 7.48% to 41.6%, depending on specific environmental conditions [6]. Conversely, improper sowing timing—whether too early or too late—can subject wheat to either premature high temperatures or late-season cold, both of which reduce growth potential and productivity [7]. For example, in Bangladesh, late sowing often exposes wheat to elevated temperatures during the grain-filling

stage, accelerating senescence and lowering yields. This issue is particularly significant in the region due to the early onset of high temperatures as the season progresses, which can severely impact yield potential. Conversely, early sowing may lead to excessive vegetative growth, increasing the risk of pest and disease infestations, especially in humid environments like those in Bangladesh [8-10]. These factors highlight the importance of optimizing sowing dates to adapt to local climate challenges and maximize wheat productivity.

In addition to optimal sowing timing, the application of plant growth regulators (PGRs) presents a promising strategy to improve wheat performance. PGRs are compounds that influence plant physiology and biochemistry, significantly contributing to growth regulation, stress tolerance, and yield enhancement [11]. Common PGRs, such as gibberellic acid (GA₃), indole-3-acetic acid (IAA), and naphthalene acetic acid (NAA), each play distinct roles in plant development. For instance, GA₃ promotes cell elongation, seed germination, and grain filling, making it a valuable tool for enhancing plant stature and yield traits [12]. In certain wheat varieties, foliar application of GA₃ has shown improvements in biological and grain yields under drought conditions, achieving increases of 12.12% and 1.47%, respectively [13]. By regulating growth parameters such as tillering, spike length, and grain weight, PGRs enable crops to better withstand environmental stressors while maximizing yield potential.

While extensive research has explored the effects of optimal sowing timing and PGR application individually, studies on their combined impact remain limited and largely inconclusive. It is plausible that integrating these

two factors could produce a synergistic effect, allowing for more robust plants that are better adapted to local environmental conditions. Aligning sowing dates with favorable weather conditions can maximize the growing season, while PGRs further bolster plant resilience and productivity by enhancing physiological growth processes. This gap in existing research underlines the need for further study to clarify the interactions between sowing timing and PGR application.

To address this gap, the present study aimed to investigate the combined effects of sowing timing and PGR application on yield traits in the wheat variety *BARI gom-33* under field conditions. By examining key parameters such as plant height, tiller number, spike length, and grain weight, grain and straw yield, this study seeks to develop insights into optimizing agronomic practices for enhanced wheat production in diverse environmental contexts.

2. MATERIALS AND METHODS

2.1 Experimental Site and Soil

The experiment was conducted at the Agronomy Field Laboratory of Bangladesh Agricultural University, Mymensingh, from November 2019 to March 2020. The experimental site is situated at 24.75°N latitude and 90.50°E longitude, with an elevation of 18 meters above sea level. The local climate is subtropical, characterized by high temperatures and heavy rainfall during the *Kharif-2* season (June to September), and scant rainfall, along with moderately low temperatures and ample sunshine, during the *Rabi* season (October to February). The soil at the experimental site belongs to the Sonatala series within the Old Brahmaputra Floodplain (AEZ-9). The detailed physiochemical characteristics of

the experimental field's soil and meteorological parameters are presented in Table 1, Table 2, and Table 3.

Table 1. Physical properties of initial soil

A. Physical Characteristics of Soil	Results
Sand (%) (0.0-0.02 mm)	20
Silt (%) (0.02-0.002 mm)	67
Clay (%) (<0.002 mm)	13
Soil textural class	Silt loam
Particle density (g/cc)	2.60
Bulk density (g/cc)	1.35
Porosity (%)	46.67

Table 2. Chemical properties of initial soil

A. Chemical Characteristics of Soil	
pH	6.80
Organic carbon (%)	1.29
Total Nitrogen (%)	0.101
Available Phosphorus (P) (ppm)	6.00
Exchangeable Potassium (K) (ppm)	6.77
Available Sulfur (S) (ppm)	10.5
Available Zinc (Zn) (ppm)	0.90
EC ($\mu\text{S cm}^{-1}$)	348

2.2 Experimental Design and Treatments

BARI gom-33 was used as the test crop. The experiment was laid out in a split-plot design with three replications. Sowing dates were assigned to the main plots, and PGRs were allocated to the sub-plots. A total of 36 plots (12 treatments \times 3 replications) were used, with a unit plot size of 2.5 m \times 2.5 m (5 m²). The planting method followed a continuous row system, with distances of 1 m between replications and 0.75 m between plots.

Table 3. Monthly record of temperature, relative humidity, rainfall, and sunshine during the period from November 2019 to March 2020 at the BAU campus

Month and Year	Air Temperature (°C)			Rainfall (mm)	Relative Humidity (%)	Sun Shine (hrs.)
	Maximum	Minimum	Average			
November 2019	29.5	18.1	23.4	1.0	81.00	204.8
December 2019	27.5	14.6	21.1	0.00	81.4	180.30
January 2020	25.08	12.85	19.40	0.00	76.74	168.9
February 2020	28.30	15.81	22.08	0.20	74.71	157.2
March 2020	27.30	18	23.30	163.7	78.52	151.6

Source: Weather Yard, department of irrigation and water management, Bangladesh Agricultural University, Mymensingh.

The experiment consisted of the following treatments:

- **Factor A: Sowing Date**
 - i. S₁: 20th November
 - ii. S₂: 5th December
 - iii. S₃: 20th December
- **Factor B: Plant Growth Regulators (PGRs)**
 - i. P₀: Water only (Control)
 - ii. P₁: Indole-3-Acetic Acid (IAA)
 - iii. P₂: Gibberellic Acid (GA₃)
 - iv. P₃: Naphthalene Acetic Acid (NAA)

2.3 Management of The Crop

The crop was planted in continuous rows, with 1 m spacing between replications and 0.75 m between plots. Sowing was carried out at a rate of 120 kg/ha, maintaining 5 cm between plants and 20 cm between rows. Recommended doses of Nitrogen (N), phosphorus (P), potassium (K), and sulfur (S) were applied at rates of 200 kg urea ha⁻¹, 160 kg triple superphosphate (TSP) ha⁻¹, 45 kg muriate of potash (MoP) ha⁻¹, and 115 kg gypsum ha⁻¹, respectively. Urea was top-dressed in three equal splits: the first at the early tillering stage (28 DAS), the second at the booting stage (46 DAS), and the third at the reproductive stage (64 DAS).

Plant growth regulators (PGRs) were sprayed at 35 and 55 days after sowing (DAS) using a sprayer (NAA @25 mg L⁻¹; GA₃ @ 15 mg L⁻¹; IAA@20 mg L⁻¹). Irrigation and intercultural operations, such as weeding, thinning, and gap-filling, were performed as needed to support optimal growth. Harvesting was conducted on three separate dates (15th March, 30th March, and 14th April) according to crop maturity, determined by the three sowing dates. For yield assessment, plants were harvested from a 1.0 m² area (1 m × 1 m) at full maturity.

2.4 Preparation of PGR Solution

To prepare a 100ppm stock solution of each plant growth regulator (PGR), the following steps were taken:

- **IAA (Indole-3-Acetic Acid):** 0.1 g of IAA powder was fully dissolved in 50 ml of 95% ethanol. Distilled water was then added to bring the total volume to 1 liter, resulting in

a 100 ppm concentration. This solution was used directly in the experiment.

- **NAA (Naphthalene Acetic Acid):** 0.1 g of NAA powder was dissolved in 1 liter of water, yielding a 100 ppm stock solution. A 40 ppm solution was then prepared from this stock by further dilution for experimental use.
- **GA₃ (Gibberellic Acid):** 0.1 g of GA₃ powder was dissolved in 1 liter of water to produce a 100 ppm stock solution, which was used directly in the experiment.

2.5 Soil Chemical Analysis

Soil texture was determined by hydrometer method as indicated by Gavlak et al. [14]. Soil pH was measured with a glass electrode pH meter in a 1:2.5 soil-to-water suspension [15]. Organic carbon was determined by the Walkley and Black wet oxidation method [16] and total nitrogen by the semi-micro Kjeldahl method [17]. Available phosphorus was extracted with 0.5 M NaHCO₃ (pH 8.5) [18] and exchangeable potassium with 1.0 N ammonium acetate (pH 7), measured via flame photometry [19]. Available sulfur was assessed using 0.15% CaCl₂ extraction [20].

2.6 Recording of Different Growth and Yield Components

Growth and yield traits, including plant height, tiller count, dry matter, spikelets per spike, and 1000-grain weight, were recorded from five randomly selected plants per plot and averaged. Leaf area was measured using an automatic leaf area meter (Type AAN-7, Hayashi DamKo Co., Japan), and leaf area index (LAI) was calculated as the ratio of total leaf area to ground area ($LAI = LA/P$), where LA represents total leaf area (cm²) and P is the ground area (cm²). Grain yield was determined from a 1 m² area at each plot's center and expressed in tons per hectare (t ha⁻¹) at 14% moisture content, while straw yield was based on sun-dried weight.

2.7 Statistical Analysis

Analysis of variance (ANOVA) was performed on all parameters using F-statistics to assess treatment effects. Means were calculated, and Tukey's HSD test at a 5% significance level was used for pairwise comparisons [21]. Data analysis was conducted by R programming language.

3. RESULTS AND DISCUSSION

3.1 Plant Growth Factors as Influenced by Sowing Date and PGRs

3.1.1 Plant height

The statistical analysis revealed significant variations in plant height due to different sowing dates and PGRs (Fig. 1). Plants sown on 20th December exhibited the greatest plant height (101.13 cm) while the lowest height was observed when plants sown on 20th November. In addition, plant height ranged from 90.21 cm in P₀ to 103.68 cm in P₂ where GA₃ was sprayed. In contrast, the maximum height was observed in the S₃:P₂ interaction (110.53 cm) while the minimum height was obtained in the control (85.00 cm).

These results aligned with those of Jamal and Mohammed [22] and Wahid et al. [23] who found that earlier sowing provided favorable environmental conditions, such as optimal temperatures, that enhanced photosynthesis and growth.

3.1.2 Leaf Area Index (LAI) as influenced by sowing date, PGRs, and their interaction

Significant effects of both sowing date and PGRs on LAI were observed (Table 4, Fig. 2). At 30, 45, 60, and 75 DAS, the highest LAI was recorded with seeds sown on 5th December and treated with GA₃, whereas the lowest LAI was observed under control conditions with water application on 20th November. These findings aligned with Liu et al. [4] who reported that timely sowing generally leads to higher LAI due to a longer vegetative growth period, allowing plants to develop more leaves and a larger canopy that enhances photosynthesis. In contrast, delayed sowing reduced LAI, as the shorter growing period limited leaf development, resulting in a smaller canopy that restricts sunlight capture and efficient photosynthesis, consistent with findings by Wahid et al. [24] and Kiss et al. [25].

Furthermore, GA₃ application significantly improved LAI, corroborating the results of Shahzad et al. [26] who found that GA₃-treated plants maintained higher LAI under stress




Fig. 1. Plant height of wheat as influenced by sowing date and PGRs. a) Plant Height affected by sowing date; b) Plant height affected by PGRs; c) Plant height affected by the interaction effect of sowing date and PGRs (Data are presented as mean \pm SEM; n = 3; $\alpha=0.05$)

(S₁= 20th November; S₂= 5th December; S₃= 20th December) (P₀= Water only; P₁= IAA; P₂= GA₃; P₃= NA)

conditions than non-treated plants. This enhancement may be attributed to GA₃'s stimulation of leaf growth, which increased leaf area and directly contributed to higher LAI, thereby improving sunlight capture and photosynthetic efficiency, as suggested by Chen et al. [27] and Shah et al. [28].

3.1.3 Dry matter accumulation of wheat as influenced by sowing date, PGRs, and their interaction

Statistical analysis showed significant variations in dry weight due to differences in sowing date, levels of plant growth regulators (PGRs), and their interaction (Table 5, Fig. 3). The highest dry matter was observed when seeds were sown on 5th December and treated with GA₃, while the lowest shoot dry weight occurred in the control treatment, where no PGRs were applied. These findings align with those of Zhiipao et al. [10] who reported that timely sowing promoted better nutrient uptake and remobilization, resulting in an 18.8% increase in post-anthesis dry matter accumulation. Liu et al. [29] similarly noted that wheat sown at an optimal time benefited from a longer growth period, allowing for greater biomass accumulation, whereas late sowing shortened the growth phases, reducing dry matter production.

Moreover, GA₃ treatment enhanced germination rates and seedling growth under both normal and osmotic stress conditions, leading to increases in shoot and root lengths as well as dry weights, as supported by Sarwar et al. [30] Guoping [31] found that GA₃ improved growth and yield performance by enhancing intercepted photosynthetically active radiation (IPAR) and

stimulating enzyme activity related to nitrogen metabolism. This increased nitrogen translocation to tillers, which contributed to better tiller development and overall dry matter accumulation [32].

3.2 Yield Contributing Characters of Wheat as Influenced by Sowing Date and PGRs

3.2.1 Yield Contributing characters of wheat as influenced by sowing date

The statistical analysis revealed significant variations in plant growth characteristics due to different sowing dates (Table 6). Plants sown on 20th December exhibited the greatest spike length (14.90 cm), and the highest number of effective spikelets (14.90), which were statistically similar to those sown on 5th December (S₂). These findings contrasted with Uddin et al. [33] who observed that wheat sown on 20th November reached the maximum plant height and spike length.

In contrast, the 20th November sowing date in the current study resulted in the highest number of effective tillers (3.16) and the greatest 1000-seed weight (48.34 g). However, crops sown on this earlier date also recorded the highest number of sterile spikelets (2.31) and the lightest grains (48 g). These results aligned with those of Jamal and Mohammed [22] and Wahid et al. [23] who found that earlier sowing provided favorable environmental conditions, such as optimal temperatures, that enhanced photosynthesis and growth, leading to greater 1000-seed weight and an increased number of effective tillers in wheat.




Fig. 2. LAI of wheat as influenced by different dates of sowing and PGRs. (Data are presented as mean; n = 3; $\alpha=0.05$)

(S₁= 20th November; S₂= 5th December; S₃= 20th December); (P₀= Water only; P₁= IAA; P₂= GA₃; P₃= NA)

Table 4. Interaction effect of date of sowing and plant growth regulators on the leaf area index of wheat

Date of sowing: Plant growth regulators	Leaf area index (LAI)			
	Days after sowing (DAS)			
	30	45	60	75
S ₁ :P ₀	0.41f	1.23e	1.30g	0.60e
S ₁ :P ₁	0.52cdef	1.56bcd	1.71cde	0.96cd
S ₁ :P ₂	0.56bcd	1.92a	1.96abc	1.33a
S ₁ :P ₃	0.44ef	1.45cde	1.50efg	0.69e
S ₂ :P ₀	0.42ef	1.29de	1.38fg	0.68e
S ₂ :P ₁	0.50cdef	1.66abc	1.74cde	1.07bc
S ₂ :P ₂	0.63ab	1.89a	2.02a	1.31a
S ₂ :P ₃	0.47def	1.45cde	1.59def	0.76de
S ₃ :P ₀	0.48def	1.24de	1.35fg	0.76de
S ₃ :P ₁	0.60abc	1.67abc	1.77bcd	1.05bc
S ₃ :P ₂	0.67a	1.84ab	2.01ab	1.23ab
S ₃ :P ₃	0.52cde	1.38cde	1.61def	0.81de
SEm (±)	0.02	0.07	0.07	0.07
Level of significance	*	*	*	*
CV (%)	10.30	10.90	8.10	13.00

Means with the same letter within the same column do not differ significantly. ** = Significant at 1% level of probability, * = Significant at 5% level of probability, NS = Not significant. (S₁= 20th November; S₂= 5th December; S₃= 20th December); (P₀= Water only; P₁= IAA; P₂= GA₃; P₃= NA)

Table 5. Interaction effect of date of sowing and plant growth regulators on the dry weight plant⁻¹ of wheat

Date of sowing: Plant growth regulators	Dry weight (g) plant ⁻¹			
	Days after sowing (DAS)			
	30	45	60	75
S ₁ :P ₀	3.66f	10.35f	17.39f	23.05j
S ₁ :P ₁	5.51cd	13.08bc	19.18cd	25.30e
S ₁ :P ₂	6.89ab	14.79a	20.48b	26.50b
S ₁ :P ₃	4.72de	11.74de	18.30e	24.34g
S ₂ :P ₀	4.31ef	12.35cd	18.81d	24.14h
S ₂ :P ₁	6.06bc	14.69a	20.82b	26.04c
S ₂ :P ₂	7.35a	15.61a	21.95a	26.87a
S ₂ :P ₃	5.72c	13.42b	19.49c	24.98f
S ₃ :P ₀	3.67f	8.90g	16.44g	23.38i
S ₃ :P ₁	4.84de	10.99ef	18.01e	25.67d
S ₃ :P ₂	6.37bc	12.97bc	19.62c	25.97c
S ₃ :P ₃	4.14ef	10.24f	17.37f	24.04h
SEm (±)	0.36	0.59	0.46	0.36
Level of significance	*	*	*	**
CV (%)	8.80	4.20	11.30	10.80

** = Significant at 1% level of probability, * = Significant at 5% level of probability, NS = Not significant. (S₁= 20th November; S₂= 5th December; S₃= 20th December)
(P₀= Water only; P₁= IAA; P₂= GA₃; P₃= NA)

3.2.2 Influence of plant growth regulators on yield contributing characters of wheat

Remarkable variations in plant growth characteristics were observed with the application of different PGRs (Table 7). Specifically, plants treated with GA₃ hormone achieved the maximum number of effective tillers

(3.89), the longest spike length (13.58 cm), the heaviest grains (49.59 g), and the highest number of effective spikelets (15.82). In contrast, the lowest values for these growth parameters were recorded under control conditions where no PGRs were applied. These results were consistent with the findings of Chen et al. [27] Lu et al. [34] Al-Tahir [35].




Fig. 3. Dry weight (DW) of wheat as influenced by sowing date and PGRs. a) DW affected by sowing date; b) DW affected by PGRs (Data are presented as mean \pm SEM; $n = 3$; $\alpha=0.05$) ($S_1=20^{\text{th}}$ November; $S_2=5^{\text{th}}$ December; $S_3=20^{\text{th}}$ December); ($P_0=$ Water only; $P_1=$ IAA; $P_2=$ GA₃; $P_3=$ NA)

Table 6. Effect of date of sowing on the yield contributing characters of wheat

Date of sowing	Effective tillers hill ⁻¹	Spike length (cm)	Effective spikelets spike ⁻¹	Sterile spikelets spike ⁻¹	1000-grain weight (g)
S ₁	3.16a	11.47b	13.26b	2.31a	48.34a
S ₂	2.98b	12.49ab	14.61a	1.56b	48.00b
S ₃	3.01b	13.38a	14.90a	1.58b	48.03b
SEm (\pm)	0.06	0.55	0.51	0.25	0.11
Level of significance	*	*	**	**	*
CV (%)	12.10	7.80	4.40	4.10	5.50

Means with the same letter within the same column do not differ significantly.

** = Significant at 1% level of probability, * = Significant at 5% level of probability, NS = Not significant. ($S_1=20^{\text{th}}$ November; $S_2=5^{\text{th}}$ December; $S_3=20^{\text{th}}$ December)

Table 7. Effect of PGRs on the yield contributing characters of wheat

Plant growth regulators	Effective tillers hill ⁻¹	Spike length (cm)	Effective spikelets spike ⁻¹	Sterile spikelets spike ⁻¹	1000-grain weight (g)
P ₀	2.32d	11.32c	12.72c	1.74	47.05d
P ₁	3.16b	12.80ab	14.73b	1.84	48.03b
P ₂	3.89a	13.58a	15.82a	1.84	49.59a
P ₃	2.82c	12.09bc	13.92b	1.85	47.80c
SEm (\pm)	0.33	0.48	0.65	0.03	0.53
Level of significance	**	**	**	NS	**
CV (%)	5.70	5.10	7.70	5.40	1.40

Means with the same letter within the same column do not differ significantly; ** = Significant at 1% level of probability, * = Significant at 5% level of probability, NS = Not significant. ($P_0=$ Water only; $P_1=$ IAA; $P_2=$ GA₃; $P_3=$ NA)

3.2.3 Interaction effect of sowing date and PGRs on the yield contributing characters of wheat

Growth and yield factors varied significantly due to the interaction between sowing date and plant growth regulators (PGRs) (Table 8). The largest spikes were recorded when plants were sown on 20th December and treated with GA₃. However, other growth parameters reached their maximum under the S₂:P₂ interaction, where GA₃ was applied in combination with the 5th December sowing date. In contrast, all growth attributes showed the lowest values when no PGRs were used, as seen in the S₁:P₀ interaction.

These results may be attributed to the optimal sowing periods, which allowed wheat to benefit from favorable environmental factors such as temperature, light, and moisture essential for both vegetative and reproductive growth. Early sowing typically led to excessive vegetative growth, making plants more susceptible to pests and diseases, while late sowing reduced the growing season, thereby limiting nutrient uptake, tillering, and grain yield, as corroborated by Kanapickas et al. [36] Tian et al. [9] and Atar [8].

Furthermore, the enhanced growth observed with GA₃ application may be attributed to its role in promoting cell division and elongation, which

resulted in taller plants with stronger stems, as supported by Sarwar et al. [30]. GA₃ treatment also appeared to improve the grain-filling process, producing heavier and more numerous grains [37].

3.2.4 Yield parameters of wheat as influenced by sowing date, PGRs, and their interaction

Significant variations in yield parameters were observed due to differences in sowing date, PGR application, and their interaction, as illustrated in Figs. 4 and 5. The highest grain yield (4.41 t ha⁻¹) was achieved when seeds were sown on 5th December (an increase of 7.59% over early sowing and 12.83% over late sowing) and treated with GA₃. However, the maximum straw yield was recorded under the S₁:P₃ treatment, where NA was applied, and seeds were sown on 20th November. In contrast, the control plot (S₁:P₀) produced the lowest yield. These findings align with Wahid et al. [24] who noted that early sowing reduced grain yield due to low temperatures during anthesis, which may have negatively affected pollen viability and thus pollination. Additionally, delaying sowing generally led to a reduction in grain yield, with yield losses of approximately 1% per day of delay due to shorter vegetative and reproductive phases, as reported by Jarecki [38] and Liu et al.

Table 8. Interaction effect of date of sowing and PGRs on the yield contributing characters of wheat

Date of sowing: Plant growth regulators	Effective tillers hill ⁻¹	Spike length (cm)	Effective spikelets spike ⁻¹	Sterile spikelets spike ⁻¹	1000-grain weight (g)
S ₁ :P ₀	2.44h	10.19g	11.58f	2.27	47.16i
S ₁ :P ₁	3.38c	11.79def	13.29cdef	2.34	48.31d
S ₁ :P ₂	3.92b	12.80bcd	15.23abcd	2.32	49.76b
S ₁ :P ₃	2.91ef	11.10fg	12.97ef	2.3	48.13e
S ₂ :P ₀	2.07i	11.32efg	13.12def	1.50	46.83j
S ₂ :P ₁	2.98de	12.81bcd	14.99abcde	1.56	47.62g
S ₂ :P ₂	4.05a	13.62abc	16.31a	1.50	50.13a
S ₂ :P ₃	2.80fg	12.22def	14.03bcde	1.67	47.40h
S ₃ :P ₀	2.47h	12.45cde	13.47cdef	1.45	47.17i
S ₃ :P ₁	3.11d	13.79ab	15.45abc	1.62	48.17de
S ₃ :P ₂	3.70b	14.33a	15.92ab	1.69	48.90c
S ₃ :P ₃	2.74g	12.94bcd	14.78abcde	1.56	47.87f
SEm (±)	0.18	0.35	0.41	0.11	0.30
Level of significance	*	*	*	NS	*
CV (%)	5.70	5.10	7.70	5.40	1.40

Means with the same letter within the same column do not differ significantly.; ** = Significant at 1% level of probability, * = Significant at 5% level of probability, NS = Not significant. (S₁= 20th November; S₂= 5th December; S₃= 20th December)(P₀= Water only; P₁= IAA; P₂= GA₃; P₃= NA)

[4]. Further supporting this, Liu et al. [6] demonstrated that optimal sowing time could maximize aboveground biomass, nonstructural carbohydrate accumulation, and canopy photosynthesis, all of which may contribute to higher yields. Similarly, Solanke et al. [39] found that optimal sowing might have enhanced germination, shoot length, and seed vigor, collectively improving grain yield.

The increased yield in GA₃-treated plants corroborated findings by Dawar et al. [40] who reported that GA₃ may have enhanced grain

filling, resulting in heavier and more numerous grains that directly contributed to higher yields. GA₃ also promoted nutrient uptake and translocation, ensuring essential nutrients were available during critical growth stages, thus enhancing plant health and productivity, as observed by Rahman et al. [41]. Additionally, GA₃ may have increased leaf area, facilitating more photosynthesis and improving grain count per spike, indirectly leading to higher yields as suggested by Farman et al. [11] and Tajdari et al. [42].




Fig. 4. Yield parameters of wheat as influenced by sowing date and PGRs. a) Grain yield affected by sowing date; b) Straw yield affected by sowing date; c) Grain yield affected by PGRs; d) Straw yield affected by PGRs (Data are presented as mean \pm SEm, n = 3; $\alpha=0.05$) (S₁= 20th November; S₂= 5th December; S₃= 20th December); (P₀= Water only; P₁= IAA; P₂= GA₃; P₃= NA)




Fig. 5. Yield parameters of wheat as influenced by interaction between sowing date and PGRs. (Data are presented as mean \pm SEm, n = 3; $\alpha=0.05$) (S₁= 20th November; S₂= 5th December; S₃= 20th December); (P₀= Water only; P₁= IAA; P₂= GA₃; P₃= NA)

4. CONCLUSION

The study demonstrated that both sowing date and PGR application especially GA₃ significantly influenced wheat growth and yield. Sowing at an optimal time provided a longer growing season, favorable weather, and reduced heat stress, while delayed sowing shortened the growth period and increased exposure to high temperatures. Furthermore, GA₃ application promoted stem elongation, increased leaf area, enhanced nutrient uptake, and improved stress tolerance. Together, these effects contributed to higher biomass, improved grain filling, and overall yield increases. Therefore, sowing on 5th December combined with the GA₃ application (S₂:P₂) could serve as an effective strategy for boosting wheat production and productivity. However, further research across diverse regions and wheat cultivars is recommended to validate these findings and develop more comprehensive guidelines for optimal GA₃ application in wheat cultivation.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGMENT

The authors are grateful to the Department of Agronomy, Bangladesh Agricultural University, Mymensingh for its support in conducting the field experiment and laboratory analyses.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Maity S, Shrivastav SP. Understanding heat stress and tolerance mechanisms in wheat (*Triticum aestivum* L.): A comprehensive review. *Journal of Advances in Biology & Biotechnology*. 2024;27(7):1196-1211.
2. Dixit AK, Sen B, Bijla S, Maiti S, Kathayat B. Contribution of wheat in global food security in changing climatic conditions: Challenges ahead and coping strategies. In *wheat science*. CRC Press. 2023;107-124.
3. Kheiralipour K, Brandão M, Holka M, Choryński A. A Review of environmental impacts of wheat production in different agrotechnical systems. *Resources*. 2024; 13(7):93.
4. Liu J, He Q, Zhou G, Song Y, Guan Y, Xiao X, Sun W, Shi, Y, Zhou K, Zhou S, Wu Y, Ma S, Wang R. Effects of sowing date variation on winter wheat yield: Conclusions for suitable sowing dates for high and stable yield. *Agronomy*. 2023;13(4).Article 4. Available:<https://doi.org/10.3390/agronomy13040991>
5. Qiao S, Harrison SP, Prentice IC, Wang H. Optimality-based modelling of wheat sowing dates globally. *Agricultural Systems*. 2023;206:103608.
6. Liu Y, Cai W, Zhu K, Xu Y, Wang W, Zhang H, Yang J. Comparison of population photosynthesis characteristics and grain yield of wheat under various sowing dates and seeding rates. *Crop Science*; 2024.
7. Gupta S. Effect of different sowing dates on growth and yield attributes of wheat in Udham Singh Nagar district of Uttarakhand, India. *Plant Archives*. 2027;17(1):232-236.
8. Atar B. Yield loss after belated sowing and effectiveness of seed vernalization in wheat. *Applied Ecology & Environmental Research*. 2024;22(2).
9. Tian Z, Yin Y, Li B, Zhong K, Liu X, Jiang D, Dai T. Optimizing planting density and nitrogen application to mitigate yield loss and improve grain quality of late-sown wheat under rice-wheat rotation; 2024.
10. Zhiipao RR, Pooniya V, Kumar D, Biswakarma N, Bainsla NK, Saikia N, Jat RD. Late-sown stress afflict post-anthesis dry matter and nutrient partitioning and their remobilization in aestivum wheat genotypes. *Journal of Agronomy and Crop Science*. 2024;210(2):e12693.
11. Farman S, Mushtaq A, Azeem MW. Plant growth regulators (PGRs) and their applications: A review. *Int. J. Chem. Biochem. Sci*. 2019;15:94-103.
12. Yuying Y, Ji L, Guo B, Li T, Guohua M, Wu K, Yang F, Zhu G, Fang L, Zeng S. Exogenous GA₃ promotes flowering in *Paphiopedilum callosum* (Orchidaceae) through bolting and lateral flower development regulation. *Horticulture Research*. 2022;9. Available:<https://doi.org/10.1093/hr/uhac09>

- 1
13. Haque MN, Pramanik SK, Hasan MA, Islam MR, Sikder S. Foliar application of potassium and gibberellic acid (GA₃) to alleviate drought stress in wheat. *Journal of Science and Technology*. 2022;1, 10.
14. Gavlak R, Horneck D, Miller R. Plant, soil, and water reference methods for the Western Region (Western Regional Extension Publication No. 125). WERA-103 Technical Committee; 2005. Available:<http://www.naptprogram.org/files/napt/western-states-method-manual-2005.pdf>
15. Michael AM. Determination of soil pH by glass electrode pH meter. *Journal of Agricultural Science*. 1965;65(2):143-145.
16. Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Science*. 1934;37(1):29-38.
17. Bremner JM, Mulvaney CS, Nitrogen-total. In A. L. Page, R. H. Miller, & D. R. Keeney (Eds.), *Methods of Soil Analysis. Part 2. Chemical and microbiological properties*. American society of agronomy, soil science society of America; 1982;595-624.
18. Olsen SR, Cole CV, Watanabe FS, Dean LA. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. U.S. Department of agriculture circular. 1954;939.
19. Knudsen D, Peterson GA, Pratt PF. Lithium, sodium, and potassium. In A. L. Page, R. H. Miller, & D. R. Keeney (Eds.), *Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties*. American Society of Agronomy, Soil Science Society of America. 1982;225-246.
20. Williams CH, Steinbergs A. Soil sulphur fractions as chemical indices of available sulphur in some Australian soils. *Australian Journal of Agricultural Research*. 1959;10(3):340-352.
21. Gomez KA. *Statistical procedures for agricultural research*. John NewYork: Wiley and Sons; 1984.
22. Jamal WM, Mohammed FHAH. Effect of hierarchy of the production of tillers in wheat (*Triticum aestivum* L.) cultivar (KM5180) under the influence of planting dates. *Research on Crops*. 2023;24(2):236-240.
23. Wahid SA, Al-Hilfy IHH. Growth and yield components of some bread wheat cultivars as affected by different sowing dates. *The Iraqi Journal of Agricultural Science*. 2018;49(2):171-178.
24. Wahid SA, Al-Hilfy IHH, Al-Abodi HMK. Effect of sowing dates on the growth and yield of different wheat cultivars and their relationship with accumulated heat units; 2017.
25. Kiss T, Balla K, Bányai J, Veisz O, Karsai I. Effect of Different Sowing Times on the Plant Developmental Parameters of Wheat (*Triticum aestivum* L.). *Cereal Research Communications*, 2014;42(2):239–251. Available:<https://doi.org/10.1556/CRC.2013.0064>
26. Shahzad K, Hussain S, Arfan M, Hussain S, Waraich EA, Zamir S, Saddique M, Rauf A, Kamal KY, Hano C. Exogenously applied gibberellic acid enhances growth and salinity stress tolerance of maize through modulating the morpho-physiological, biochemical and molecular attributes. *Biomolecules*. 2021;11(7):1005. Available:<https://doi.org/10.3390/biom11071005>
27. Chen L, Hao L, Condon AG, Hu YG. Exogenous GA₃ application can compensate the morphogenetic effects of the GA-responsive dwarfing gene Rht12 in Bread Wheat. *Plos One*, 2014;9(1):e86431. Available:<https://doi.org/10.1371/journal.pone.0086431>
28. Shah SH, Islam S, Mohammad F, Siddiqui MH. Gibberellic acid: A versatile regulator of plant growth, development and stress responses. *Journal of Plant Growth Regulation*. 2023;42(12): 7352–7373. Available:<https://doi.org/10.1007/s00344-023-11035-7>
29. Liu K, Zhang C, Guan B, Yang R, Liu K, Wang Z, Wang X. The effect of different sowing dates on dry matter and nitrogen dynamics for winter wheat: an experimental simulation study. *PeerJ*. 2021;9:e11700.
30. Sarwar G, Anwar T, Malik M, Rehman H UR, Danish S, Alahmadi TA, Ansari MJ. Evaluation of potassium-enriched biochar and GA₃ effectiveness for Improving wheat growth under drought stress. *BMC Plant Biology*. 2023;23(1):615. Available:<https://doi.org/10.1186/S12870-023-04613-0>
31. Guoping Z. Gibberellic Acid₃ modifies some growth and physiologic effects of

- paclobutrazol (PP₃₃) on wheat. Journal of Plant Growth Regulation. 1997;16(1):21–25.
Available:<https://doi.org/10.1007/PL00006970>
32. Wang Y, Ren T, Lu J, Cong R, Hou W, Liu T, Hussain S, Li X. Exogenously applied gibberellic acid improves the growth and yield performance of inferior rice tillers grown under different nitrogen levels. Acta Physiologiae Plantarum. 2016;39(1), 5.
Available:<https://doi.org/10.1007/S1738-016-2307-3>
 33. Uddin R, Islam M, Ullah M, Hore P, Paul S. Grain growth and yield of wheat as influenced by variety and sowing date. Bangladesh Agronomy Journal. 2016;18(2):97–104.
Available:<https://doi.org/10.3329/baj.v18i2.28911>
 34. Lu Q, Lu S, Wang M, Cui C, Condon AG, Jatayev S, Hu YG. The exogenous GA₃ greatly affected the grain-filling process of semi-dwarf gene Rht4 in bread wheat. Physiologia Plantarum. 2022;174(3):e13725.
 35. Al-Tahir FM. Relationship between tiller development and productivity with yield and its components for wheat and oat under the influence of All and GA₃. European Academic Research. 2014;9(2):11466-11485.
 36. Kanapickas A, Vagusevičienė I, Sujetovienė G. The effects of different sowing dates on the autumn development and yield of winter wheat in Central Lithuania. Atmosphere. 2024;15(6):738.
 37. Anwar T, Shehzadi A, Qureshi H, Shah MN, Danish S, Salmen SH, Ansari MJ. Alleviation of cadmium and drought stress in wheat by improving growth and chlorophyll contents amended with GA₃ enriched deashed biochar. Scientific Reports. 2023;13(1):18503.
 38. Jarecki W. Response of winter wheat to delayed sowing and varied nitrogen fertilization. Agriculture. 2024;14(1), Article 1.
Available:<https://doi.org/10.3390/agriculture14010121>
 39. Solanke AP, Raut D, Kalbhor SU, Nalbale SR, Lande SS. Impact of varied sowing dates on seed quality parameters in wheat. Journal of Advances in Biology & Biotechnology. 2024;27(9):116-122.
 40. Dawar K, Rahman U, Alam SS, Tariq M, Khan A, Fahad S, Datta R, Danish S, Saud S, Noor M. Nitrification inhibitor and plant growth regulators improve wheat yield and nitrogen use efficiency. Journal of Plant Growth Regulation. 2022;41(1):216–226.
Available:<https://doi.org/10.1007/s00344-020-10295-x>
 41. Rahman HMA, Ahmed BEAM, Ahmed MO. Effect of plant growth regulators (GA₃, NAA) and water stress on leaf area and yield attributes of wheat. International Journal of Innovative Science and Research Technology. 2018;3(5):1–5.
 42. Tajdari HR, Soleymani A, Montajabi N, Naderi Darbaghshahi MR, Javanmard HR. The effect of foliar application of plant growth regulators on functional and qualitative characteristics of wheat (*Triticum aestivum* L.) under salinity and drought stress conditions. Applied Water Science. 2024;14(6):126.
Available:<https://doi.org/10.1007/S13201-024-02203-5>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/126457>